

Importation of Fresh Garden Bean, *Phaseolus vulgaris* Linnaeus, from Egypt into the Continental United States

A Qualitative, Pathway-Initiated Pest Risk Assessment

December 11, 2012

Version 1

Agency Contact:

Plant Epidemiology and Risk Analysis Laboratory
Center for Plant Health Science and Technology

United States Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine
1730 Varsity Drive, Suite 300
Raleigh, NC 27606

Executive Summary

The Animal and Plant Health Inspection Service (APHIS) of the United States Department of Agriculture (USDA) prepared this risk assessment document to examine plant pest risks associated with importing commercially produced fresh garden beans, *Phaseolus vulgaris* (Fabaceae), for consumption, in pods or shelled, from Egypt into the continental United States. As the market access request by Egypt did not specify any production, post-harvest, or transportation practices, we assumed no such practices when preparing this risk assessment.

Based on the scientific literature, port-of-entry pest interception data, and information from the government of Egypt, we developed a list of all potential pests with actionable regulatory status for the continental United States that are known to occur in Egypt and to be associated with garden beans anywhere in the world. From this list, we identified seven organisms that have a reasonable likelihood of being on garden beans at the time of harvest and remaining throughout harvest. We analyzed these seven pests in more detail.

Of the pests selected for further analysis, we determined that the following are *not* candidates for risk management, either because there is no endangered area within the continental United States, they did not meet the threshold to likely cause unacceptable consequences of introduction, or because they received a Negligible overall risk rating for likelihood of introduction (i.e., entry plus establishment) into the endangered area via the import pathway: *Bruchus tristis* Boheman (Coleoptera: Bruchidae), *Icerya seychellarum* (Westwood) (Hemiptera: Monophlebidae), and *Maconellicoccus hirsutus* (Green) (Hemiptera: Pseudococcidae).

We determined that the following arthropod pests are candidates for risk management, because they **met the threshold to likely cause unacceptable consequences of introduction**, and they received an overall **likelihood of introduction** risk rating **above** Negligible:

Taxonomy	Scientific Name	Likelihood of Introduction overall rating
Lepidoptera: Lycaenidae	<i>Lampides boeticus</i> L.	Medium
Lepidoptera: Noctuidae	<i>Chrysodeixis chalcides</i> (Esper)	Medium
	<i>Helicoverpa armigera</i> Hübner	Medium
	<i>Spodoptera littoralis</i> (Boisduvalle)	Medium

Detailed examination and choice of appropriate phytosanitary measures to mitigate pest risk are part of the pest risk management phase within APHIS and are not addressed in this document.

Table of Contents

Executive Summary	2
1. Introduction	4
1.1. Background	4
1.2. Initiating event.....	4
1.3. Determination of the necessity of a weed risk assessment for the commodity	4
1.4. Description of the pathway.....	5
2. Pest List and Pest Categorization	6
2.1. Pests considered but not included on the pest list	6
2.2. Pest list	9
2.3. Pests selected for further analysis	18
3. Assessing Pest Risk Potential	18
3.1. Introduction	18
3.2. Assessment results.....	19
4. Summary and Conclusions of Risk Assessment.....	33
5. Acknowledgements	34
6. Literature Cited	35
7. Appendix.....	43
Appendix A. Pests with non-actionable regulatory status.....	43

1. Introduction

1.1. Background

This document was prepared by the Plant Epidemiology and Risk Analysis Laboratory of the Center for Plant Health Science and Technology, USDA Animal and Plant Health Inspection Service (APHIS), Plant Protection and Quarantine (PPQ), to evaluate the risks associated with the importation of commercially produced fresh garden beans, *Phaseolus vulgaris* Linnaeus, for consumption, in pods or shelled, from Egypt into the continental United States.

The International Plant Protection Convention (IPPC) provides guidance for conducting pest risk analyses. The methods used here are consistent with guidelines provided by the IPPC, specifically the International Standard for Phytosanitary Measures (ISPM) on ‘Pest Risk Analysis for Quarantine Pests, Including Analysis of Environmental Risks and Living Modified Organisms’ (IPPC, 2011). The use of biological and phytosanitary terms is consistent with the ‘Glossary of Phytosanitary Terms’ (IPPC, 2012).

Three stages of pest risk analysis are described in international standards: Stage 1, Initiation; Stage 2, Risk Assessment; and Stage 3, Risk Management. This document satisfies the requirements of Stages 1 and 2.

This is a qualitative risk assessment. We express the risk based on qualitative ratings for the likelihood and consequences of pest introduction via the imported green beans from the Egypt. The details of the methodology and rating criteria are found in the *Guidelines for Plant Pest Risk Assessment of Imported Fruit and Vegetable Commodities, Version 6.0* (PPQ, 2012).

The appropriate risk management strategy for a particular pest depends on the risk posed by that pest. Identification of appropriate phytosanitary measures to mitigate pest risk is undertaken in Stage 3 (Risk Management) and is not covered in this risk assessment. Risk management will be specified in a separate document.

1.2. Initiating event

The importation of fruits and vegetables for consumption into the United States is regulated under Title 7 of the Code of Federal Regulations, Part 319.56 (7 CFR §319.56, 2012). Currently, under this regulation, the entry of green beans for consumption in pods or shelled from Egypt into the continental United States is not authorized. This assessment was prepared in response to a request Dr. Ali Soliman, Head of the Central Administration for Plant Quarantine, Ministry of Agriculture and Land Reclamation, Cairo, Egypt to change the Federal Regulation to allow entry (Soliman, 2007).

1.3. Determination of the necessity of a weed risk assessment for the commodity

In some cases, an imported commodity may have the potential of becoming invasive in the PRA area. The likelihood that this may happen is evaluated in a weed risk assessment, conducted separately from the commodity risk assessment.

Weed risk assessments do not need to be conducted for plant species that are widely established (native or naturalized) or cultivated in the PRA area, for commodities that are already enterable into the PRA area from other countries, or when the plant part(s) cannot easily propagate on their own or be propagated. We determined that a weed risk assessment is not needed for *Phaseolus vulgaris* because it is widely cultivated in the United States, and is enterable from other countries (APHIS, 2012).

1.4. Description of the pathway

The IPPC (2011) defines a pathway as “any means that allows the entry or spread of a pest.” In the context of commodity pest risk assessments, the *pathway* is the commodity to be imported, together with all the processes the commodity undergoes that may have an impact on pest risk. In this risk assessment, the specific pathway of concern is the importation of fresh garden beans (*Phaseolus vulgaris* L.) for consumption, in pods or shelled, from Egypt into the continental United States; the movement of this commodity provides a potential pathway for the introduction and/or spread of plant pests.

The following description of this pathway focuses on the conditions that may affect plant pest risk, including morphological and physiological characteristics of the commodity, as well processes the commodity will undergo from production in Egypt through importation and distribution in the continental United States. These conditions provided the basis for creating the pest list and assessing the likelihood of introduction of the pests selected for further analysis; therefore, all components of the pathway, as they are described below, should be considered mandatory conditions for importation of the commodity.

1.4.1. Description of the Commodity

Egypt will be exporting fresh garden beans for consumption, in pods (Fig. 1) or shelled (Fig. 2). The beans are harvested before the seeds mature and harden (Lerner, 2001). This means that the seeds of fresh garden beans, whether they are left inside the pods or shelled, are not mature and therefore cannot produce new plants. Consequently, there is no concern that the seeds may be diverted for planting. Furthermore, the seeds are not a viable pathway for pathogens that are exclusively seed-transmitted or for arthropods that require the mature seed to complete their development. If unrefrigerated, fresh garden beans decompose within a few days, thus ceasing to provide suitable conditions for internal pests.



Figure 1. Fresh garden beans in pods. (Image by J.R. Manhart, http://www.metafro.be/prelude/view_plant?pi=09910)

1.4.2. Production and harvest procedures in the exporting area

Production and harvesting procedures have not been specified by the exporting country and are therefore not being considered as part of the risk assessment.

1.4.3. Post-harvest procedures in the exporting area

Post-harvest procedures have not been specified by the exporting country and are therefore not being considered as part of the risk assessment.

1.4.4. Shipping and storage conditions

Beans will be shipped in 3 to 5 kg carton boxes throughout the year via air and maritime shipments (Ministry of Agriculture and Land Reclamation, 2011). Based on standard industry practices, we assume that the beans will be shipped at temperatures of 4-7 degrees C (McGregor, 1987).

2. Pest List and Pest Categorization

In this section, we identify the plant pests with actionable regulatory status for the continental United States that could potentially become established in the continental United States as a result of the importation of fresh green beans (*Phaseolus vulgaris* Linnaeus) for consumption, in pods or shelled, from Egypt, and we determine which of these pests meet the criteria for further analysis (Table 2). Pests are considered to be of regulatory significance if they are actionable at U.S. ports-of-entry. Actionable pests include quarantine pests, pests considered for or under official control, and pests that require evaluation for regulatory action.

2.1. Pests considered but not included on the pest list

2.1.1. Pests with weak evidence for association with the commodity or for presence in the export area

***Bactrocera cucurbitae*.** De Meyer et al. (2012) report that extensive surveys in Egypt have shown that *B. cucurbitae* does not occur there and that an old report of *B. cucurbitae* from the “Lower Nile Valley, Egypt” could not be confirmed or traced to its origin. EPPO considers this pest to be absent from Egypt (EPPO, 2012a).

***Ceratitis capitata*.** Although Thomas et al. (2005) listed *P. vulgaris* as a *C. capitata* host of “unknown importance,” we found no substantiating evidence that *C. capitata* commonly feeds on *P. vulgaris*. For example, during 37 years of surveys in Hawaii this fruit fly has never been detected on *P. vulgaris* (PERAL, 2008, Liquido et al., 1990; Liquido et al., 1991; Thomas et al., 2005). Therefore, we concluded there was insufficient evidence of host association.

***Cryptoblabes gnidiella*.** We think all references to this pest feeding on beans can be traced back to a single sentence (Mau and Kessing, 1992): “In Hawaii, this pest has been recorded on the following hosts: Christmas berry, coffee, corn, green beans, sorghum”. We found no other credible evidence to corroborate beans as a host of *C. gnidiella*.

***Dacus ciliatus*.** CABI (2012) states “there are a few reports from hosts other than Cucurbitaceae, namely *Adenia gummifera* (Passifloraceae), *Gossypium* sp. (Malvaceae), *Solanum lycopersicum* (Solanaceae) and *Phaseolus* sp. (Fabaceae), but these are not common hosts and may represent

aberrant associations or a confused host range”. We found no other credible evidence that *D. ciliatus* feeds on *P. vulgaris* or any plant in the family Fabaceae, so we did not list it.

Retithrips syriacus. While *Phaseolus vulgaris* is listed as a host in CABI (2012), we found no other evidence that this thrips feeds on beans.

Tuta absoluta. All mentions in the literature of this pest feeding on beans can be traced back to a single report by the NPPO of Italy. The only information available regarding this report is “The NPPO of Italy has recently informed the EPPO Secretariat that *Tuta absoluta* (Lepidoptera: Gelechiidae – EPPO A1 List) has been found on *Phaseolus vulgaris* plants in Sicilia. So far, the pest had mainly been observed on tomato crops (*Lycopersicon esculentum*)” (EPPO, 2009). In the absence of further details and given that *T. absoluta* is a well-researched pest and has never been reported on beans anywhere else in the world it occurs, we believe that this is insufficient evidence that *P. vulgaris* is a host of *T. absoluta*.

***Ralstonia solanacearum* Race 3 Biovar 2 (RSR3B2)**. Conflicting evidence exists about the host status of *P. vulgaris*. Citing previous reports, Janse et al. (1994) suggested that *P. vulgaris* was a host, despite the fact that the pathogenicity tests of *R. solanacearum* on field-grown mature field beans were not performed. The most current information (Álvarez et al., 2007; Allen, 2012; Allen, 2012) suggests that *P. vulgaris* is a non-host, which is what we concluded.

2.1.2. Organisms with non-actionable regulatory status

We found some evidence of the organisms listed in Appendix A being associated with garden beans for consumption, in pods or shelled and being present in the Egypt; however, because these organisms have non-actionable regulatory status for the continental United States, we did not include them below (Table 2) in this risk assessment.

2.1.3. Organisms identified only to the genus level

In commodity import risk assessments, the taxonomic unit for pests selected for evaluation beyond the pest categorization stage is usually the species (IPPC, 2011), as assessments focus on organisms for which biological information is available. Therefore, generally, we do not assess risk for organisms identified only to the genus level, in particular if the genus in question is reported in the import area. Often there are many species within a genus, and we cannot know if the unidentified species occurs in the import area and, consequently, whether it has actionable regulatory status for the import area. On the other hand, if the genus in question is absent from the import area, any unidentified organisms in the genus can have actionable status; however, because such an organism has not been fully identified, we cannot properly analyze its likelihood and consequences of introduction.

In light of these issues, we usually do not include organisms identified only to the genus level in the main pest list. Instead, we address them separately in this sub-section (Table 1). The information here can be used by risk managers to determine if measures beyond those intended to mitigate fully identified pests are warranted. Often, however, the development of detailed assessments for known pests that inhabit a variety of ecological niches, such as internal fruit feeders or foliage pests, allows effective mitigation measures to eliminate the known organisms as well as similar but incompletely identified organisms that inhabit the same niche.

Table 1. Organisms identified to the genus level that are reported on *Phaseolus vulgaris* in Egypt and that have actionable or undetermined regulatory status.

Pest Name	Evidence of presence on <i>Phaseolus vulgaris</i> in Egypt	Genus present in the continental United States?	Regulatory status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
Acari: Tetranychidae						
<i>Tetranychus</i> spp.	CABI, 2012; Seif et al., 2001	Yes (CABI, 2012)	U	Leaves (Seif et al., 2001)	No	
Coleoptera: Meloidae						
<i>Coryna</i> spp.	Seif et al., 2001	No (Arnett et al., 2002)	A	Flowers (Seif et al., 2001)	No	
<i>Mylabris</i> spp.	Seif et al., 2001	No (<i>Mylabris</i> sp. in the family Meloidae are not listed as present) (Arnett et al., 2002)	A	Flowers (Seif et al., 2001)	No	
Collembola: Entomobryidae						
<i>Seira</i> sp.	Harakly and Assem, 1978	Yes (based on <i>S. punctata</i>) (Bellinger et al., 2012)	U	Occurs in soil (Muturi et al., 2009)	No	
Diptera: Agromyzidae						
<i>Liriomyza</i> sp.	Elbadry et al., 2006	Yes (based on <i>L. huidobrensis</i>) (CABI, 2012)	U	Leaves (Elbadry et al., 2006)	No	
Hemiptera: Aphididae						
<i>Aphis</i> spp.	Farrag and Zakzouk, 2000	Yes (CABI, 2012)	U	Flowers, leaves, shoots, entire plant (based on data for <i>A. fabae</i>) (CABI, 2012)	No	This external feeder (CABI, 2012) is highly unlikely to remain after harvesting.
Hemiptera: Cicadellidae						
<i>Empoasca</i> spp.	Karel and Autrique, 1989	Yes (based on <i>E. fabae</i>) (CABI, 2012)	U	Leaves (based on <i>E. fabae</i>) (CABI, 2012)	No	
Lepidoptera: Noctuidae						
<i>Agrotis</i> spp.	CABI, 2012; Seif et al., 2001	CABI, 2012	U	Girdles plants at soil level or below, indicating stems are attacked (Seif et al., 2001)	No	

Pest Name	Evidence of presence on <i>Phaseolus vulgaris</i> in Egypt	Genus present in the continental United States?	Regulatory status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
<i>Spodoptera</i> spp.	CABI, 2012; Seif et al., 2001	CABI, 2012	U	Girdles plants at soil level or below, indicating stems are attacked (Seif et al., 2001)	No	

^aA=Actionable, U=Undetermined. If the genus does not occur in the continental United States, the organism has actionable status. If the genus occurs in the continental United States, the organism has undetermined regulatory status, because we cannot know if the unidentified species is one that occurs in the continental United States.

^bThe plant part(s) listed are those for the plant species under analysis. If the information is extrapolated, such as from plant part association on other plant species, this is noted.

^c“Yes” indicates the pest has a reasonable likelihood of being associated with the harvested plant part(s).

2.2. Pest list

Below, we list the actionable pests associated with fresh green beans for consumption, in pods or shelled, that occur in Egypt (Table 2). The list comprises those actionable pests that occur in Egypt on any host and are reported to be associated with fresh green beans for consumption, in pods or shelled, whether in Egypt or elsewhere in the world. For each pest, we indicate 1) the part of the imported plant species with which the pest is generally associated, and 2) whether the pest has a reasonable likelihood of being associated, in viable form, with the commodity following harvesting from the field and prior to any post-harvest processing. We developed this pest list based on the scientific literature, port-of-entry pest interception data, and information provided by the government of Egypt. Pests in shaded rows are pests identified for further evaluation, as we consider them reasonably likely to be associated with the harvested commodity; we summarize these pests in a separate table (Table 3).

Table 2. Actionable pests reported on *Phaseolus vulgaris* (in any country) and present in Egypt (on any host)

Pest Name	Evidence of presence in Egypt	Host status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
Acari: Tetranychidae					
<i>Eutetranychus africanus</i> (Tucker)	El Kifl et al., 1974	Type 1 (El Kifl et al., 1974)	Leaves (based upon <i>E. orientalis</i> below)	No	
<i>Eutetranychus orientalis</i> Klein	El Kifl et al., 1974	Type 1 (Hill, 1983; Migeon and Dorkeld, 2006; El Kifl et al., 1974)	Leaves [based on information for soybeans (Sullivan and Jones, 2010)]	No	Present in Hawaii (Heu, 2007).
Coleoptera: Bruchidae					
<i>Bruchus tristis</i> Boheman	Gentry, 1965	Type 1 (Gentry, 1965)	Seed (Gentry, 1965)	Yes	Larvae develop in seeds (Kergoat et al., 2004).
Coleoptera: Chrysomelidae					
<i>Aulacophora foveicollis</i> Lucas (= <i>Rhaphidopalpa foveicollis</i> (Lucas)) (CABI, 2012)	CABI, 2012	Type 1 (CABI, 2012)	Fruit, flowers, leaves, roots, stems (CABI, 2012)	No. Adults feed externally and are very mobile; they are highly unlikely to remain after harvest.	Evidence that this species feeds on <i>P. vulgaris</i> is very weak.
Coleoptera: Scarabaeidae					
<i>Tropinota squalida</i> (Scopoli)	Gentry, 1965	Type 1 (Gentry, 1965)	Flowers (Gentry, 1965)	No	
Diptera: Agromyzidae					
<i>Chromatomyia horticola</i> Goureau (= <i>Phytomyza horticola</i> Goureau; <i>P. atricornis</i> (partim.) Meigen) (CABI, 2012)	CABI, 2012; Hammad, 1978	Type 1, <i>Phaseolus</i> (beans) are a host (CABI, 2012)	Leaves (CABI, 2012)	No	
<i>Liriomyza bryoniae</i> Kaltenbach	EPPO, 2006b	Type 1 (CABI, 2012; Harakly and Assem, 1978)	Leaves, stems (CABI, 2012; Spencer, 1965)	No	

Pest Name	Evidence of presence in Egypt	Host status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
<i>Liriomyza congesta</i> (Becker)	Hammad, 1978	Type 1 (El Kifl et al., 1974)	Leaves (Al-Azawi, 1967)	No	
<i>Melanagromyza sojae</i> (Zehntner)	Elbadry et al., 2006	Type 1 (Elbadry et al., 2006)	Leaves, stems (CABI, 2012; Spencer, 1973)	No	
<i>Ophiomyia phaseoli</i> Tryon (= <i>Melanagromyza phaseoli</i> Vanschuytebroeck) (CABI, 2012)	PPQ, 2002	Type 1 (PPQ, 2002)	Stems, leaves (Hill, 1983; Spencer, 1973)	No	Present in Hawaii (PPQ, 2002).
Hemiptera: Aleyrodidae					
<i>Bemisia afer</i> (Priesner & Hosny)	Abd-Rabou and Ahmed, 2008	Type 1 (Thindwa and Khonje, 2005)	Leaves (EPPO, 2004)	No	
Hemiptera: Cicadellidae¹					
<i>Balclutha hebe</i> (Kirkaldy)	Karel and Autrique, 1989	Type 1 (reported on beans) (Karel and Autrique, 1989)	Leaves (Karel and Autrique, 1989)	No	
<i>Balclutha rosea</i> (Scott)	Karel and Autrique, 1989	Type 1 (reported on beans) (Karel and Autrique, 1989)	Leaves (Karel and Autrique, 1989)	No	
<i>Balclutha saltuella</i> (Kirschbaum)	Karel and Autrique, 1989	Type 1 (reported on beans) (Karel and Autrique, 1989)	Leaves (Karel and Autrique, 1989)	No	
<i>Cicadulina chinai</i> (Ghaui)	Hashem et al., 2009	Type 1 (Hashem et al., 2009)	Leaves (based upon barley, maize and wheat) (Ammar et al., 1989)	No	
<i>Empoasca decedens</i> (Paoli) (= <i>Asymmetrasca decedens</i> (Paoli)) (Jacas et al., 1997)	Hashem et al., 2009	Type 1 (Hashem et al., 2009)	Leaves, shoots (based upon almond) (Jacas et al., 1997)	No	

¹ Most pests in this family are very mobile, and therefore highly unlikely to remain after harvest.

Pest Name	Evidence of presence in Egypt	Host status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
<i>Empoasca decipiens</i> Paoli	Raupach et al., 2002	Type 1 (Avidov and Harpaz, 1969)	Leaves (Avidov and Harpaz, 1969)	No	
<i>Empoasca distinguenda</i> Paoli	El Kifl et al., 1974	Type 1 (El Kifl et al., 1974)	Leaves, fruits (based upon green pepper and castor oil plants) (El-Dessouki and Hosny, 1969)	No. This very mobile external feeder (El-Dessouki and Hosny, 1969) is highly unlikely to remain after harvesting.	
<i>Empoasca lybica</i> (de Bergevin) (= <i>E. signata</i> (Haupt); <i>Jacobiasca lybica</i> (de Bergevin)) (CABI, 2012)	Khalafallah et al., 2006	Type 1 (Abate and Ampofo, 1996)	Leaves (Avidov and Harpaz, 1969)	No	
<i>Neolimnus egyptiacus</i> Matsumura (Membracoidea of the World Database, 2010) ²	Karel and Autrique, 1989; Membracoidea of the World Database, 2010	Type 1 (reported on beans) (Karel and Autrique, 1989)	Leaves (Karel and Autrique, 1989)	No	
<i>Orosius albicinctus</i> (= <i>O. orientalis</i> Matsumura) (CABI, 2012)	Khodeir, 2006	Type 1 (Khodeir, 2006)	Leaves (CABI, 2012)	No	
<i>Trialeurodes ricini</i> (Misra)	EPPO, 2000	Type 1 (Thindwa and Khonje, 2005)	Fruit, leaves (EPPO, 2000; Shishehbor and Brennan, 1995)	No. Cicadellidae are very mobile external feeders, and are unlikely to remain throughout harvesting.	

² We assumed this was misspelled as *N. aegyptiacus* (Karel and Autrique, 1989), and *Heolimnus aegyptiacus* (Mats.) (El Kifl et al., 1974)

Pest Name	Evidence of presence in Egypt	Host status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
Hemiptera: Lygaeidae					
<i>Spilostethus longulus</i> Dallas	Gentry, 1965	Type 1 (Gentry, 1965)	Leaves, seeds, stems (based upon <i>S. pandurus</i> on <i>Calotrcpis</i>) (Schaefer and Panizzi, 2000)	No. This very mobile external feeder (based upon Roselle; Abdel-Moniem et al., 2011) is highly unlikely to remain after harvest.	
Hemiptera: Miridae					
<i>Trigonotylus brevipes</i> Jakowlef	Gentry, 1965	Type 1 (Gentry, 1965)	Leaves, flowers (based on <i>T. caelestialium</i> on Italian rye grass) (Shiba et al., 2011)	No	Present in Guam and Samoa (Unsinger, 1951).
Hemiptera: Monophlebidae					
<i>Icerya seychellarum</i> (Westwood)	CABI/EPPO, 2008	Type 1 (Williams and Watson, 1990)	Flowers, fruit, leaves, stems (PPQ, 2002)	Yes	
Hemiptera: Pentatomidae					
<i>Agonoscelis puberula</i>	Gentry, 1965	Type 1 (Gentry, 1965)	Unknown, but assumed to be leaves, stems and possibly pods, based on general feeding habits of Pentatomidae (Triplehorn et al., 2005).	No. This very mobile, external pest (Triplehorn et al., 2005) is highly unlikely to remain after harvesting.	Present in Texas, New Mexico, and Arizona (Thomas et al., 2003) and not under official control.
Hemiptera: Pseudococcidae					
<i>Maconellicoccus hirsutus</i> (Green) (= <i>Phenacoccus hirsutus</i> Green) (CABI, 2012)	CABI/EPPO, 1997	Type 1 (Ben-Dov et al., 2012)	Fruit, leaves, stems, whole plant (Caribbean Pest Information Network, No Date)	Yes	Present in California, Florida, and Texas (CABI, 2012).
Lepidoptera: Lycaenidae					
<i>Lampides boeticus</i> Linnaeus (= <i>Cosmolyce baeticus</i> L.) (El Kifl et al., 1974; Williams, 2008)	Commonwealth Institute of Entomology, 1984	Type 1 (Grund, 2002)	Flowers, pods (Zimmerman, 1958)	Yes	Present in Hawaii (Zimmerman, 1958).

Pest Name	Evidence of presence in Egypt	Host status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
Lepidoptera: Noctuidae					
<i>Agrotis segetum</i> (Denis & Schiffermüller)	C.A.B. International, 1987	Type 1 (Seif et al., 2001)	Stems (Seif et al., 2001)	No	
<i>Autographa gamma</i> (Linnaeus) (= <i>Plusia gamma</i> (Linnaeus) (CABI, 2012)	PPQ, 2002	Type 1 (PPQ, 2002)	Fruit, flowers, leaves (Sullivan and Jones, 2010)	No. This very mobile external feeder (CABI, 2012; PPQ, 2002) is highly unlikely to remain after harvesting.	
<i>Chrysodeixis chalcites</i> (Esper) (= <i>Plusia chalcites</i> Esper) (CABI, 2012)	Commonwealth Institute of Entomology, 1977	Type 1 (Robinson et al., 2011)	Leaves, pods (Sullivan and Jones, 2010)	Yes	Present in Hawaii (Commonwealth Institute of Entomology, 1977).
<i>Helicoverpa armigera</i> Hübner (= <i>Heliothis armigera</i> Hübner) (CABI, 2012)	International Institute of Entomology, 1993	Type 1 (Abate and Ampofo, 1996; Mansour et al., 1981)	Flowers, Flower buds, leaves, pods (Ampofo, 1994)	Yes	
<i>Spodoptera exempta</i> Walker	CABI, 2012	Type 1 (Robinson et al., 2007a)	Leaves, shoots, stems (CABI, 2012)	No	Present in California, Hawaii, Kansas, Oregon, Washington, and Wisconsin (CABI, 2012).
<i>Spodoptera littoralis</i> (Boisduval) (= <i>Prodenia litura</i> Fabricius sensu auctorum) (CABI, 2012)	PPQ, 2002	Type 1 (Afifi and El-Whab, 1990; Robinson et al., 2007b)	Flowers, leaves, pods/seeds, stems (based upon soybeans) (Sullivan and Jones, 2010)	Yes	<i>S. littoralis</i> feeds internally on fruit (CABI, 2012)
<i>Syngrapha circumflexa</i> (L.) (= <i>Cornutiplusia circumflexa</i> (L.)) (Hächler, 1986; Harakly, 1975)	El Kifl et al., 1974; Harakly, 1975	Type 1, reported on bean (El Kifl et al., 1974) and <i>Phaseolus</i> (Harakly, 1975)	Leaves (based on <i>Phaseolus</i> spp.) (Harakly, 1975)	No	

Pest Name	Evidence of presence in Egypt	Host status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
<i>Thysanoplusia orichalcea</i> (Fabricius) (= <i>Diachrysia orichalcea</i> (Fabricius)) (CABI, 2012)	Zhang, 1994	Type 1 (CABI, 2012)	Fruit, leaves (CABI, 2012)	No. This very mobile external feeder (CABI, 2012) is highly unlikely to remain after harvesting.	
Lepidoptera: Sphingidae					
<i>Agrius convolvuli</i> (Linnaeus) (= <i>Herse convolvuli</i> (Linnaeus)) (CABI, 2012)	EcoPort, 2012	Type 1 (reported on <i>Phaseolus</i> spp. and beans) (Hill, 1983)	Flowers, leaves (Hill, 1983)	No	Present in Hawaii (EcoPort, 2012).
Orthoptera: Acrididae					
<i>Aiolopus strepens</i> Latr.	El Kifl et al., 1974	Type 1 (El Kifl et al., 1974)	Leaves, whole plant (extrapolated from <i>A. simulatrix</i>) (Anonymous, 1978)	No	This very mobile external feeder (El-Minshawy and El-Hinnawy, 1976) is highly unlikely to remain after harvest.
<i>Diaboloecatantops axillaris</i> (Thunberg)	CABI, 2012	Type 1 (reported on <i>Phaseolus</i>) (CABI, 2012)	Leaves, pods, stems (CABI, 2012)	No. This pest is a very mobile external feeder (CABI, 2012). See remarks for <i>Aiolopus strepens</i> .	
<i>Eyprepocnemis plorans plorans</i> (Charpentier) (= <i>Euprepocnemis plorans</i> (Charpentier) (Bisby et al., 2012)	El Kifl et al., 1974	El Kifl et al., 1974	Leaves (based upon tobacco and broad beans) (Ascher et al., 1989; Rungs, 1962)	No	
<i>Locusta migratoria</i> (Linnaeus)	CABI, 2012	Type 1 (reported on <i>Phaseolus</i>) (CABI, 2012)	Leaves, pods, stems (CABI, 2012)	No. This pest is a very mobile external feeder (CABI, 2012). See remarks for <i>Aiolopus strepens</i> .	

Pest Name	Evidence of presence in Egypt	Host status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
<i>Schistocerca gregaria</i> (Forskål)	EcoPort, 2012	Type 1 (<i>Phaseolus</i> sp. listed as a host) (van Huis et al., 2008)	Flowers, fruit, leaves, shoots, stems, seeds, entire plant (CABI, 2012)	No. This pest is a very mobile external feeder (CABI, 2012). See remarks for <i>Aiolopus strepens</i> .	Highly polyphagous pest (CABI, 2012).
Orthoptera: Pyrgomorphidae					
<i>Chrotogonus lugubris</i> Blanch.	El Kifl et al., 1974	Type 1 (El Kifl et al., 1974)	Leaves [extrapolated from clover, cotton, bean (<i>Vicia faba</i>), and wheat] (Abdel Rahman, 2001)	No	This pest is a very mobile external feeder (Abdel Rahman, 2001).
Nematodes					
<i>Heterodera cajani</i>	CABI, 2012	Type 1 (CABI, 2012)	Root (CABI, 2012)	No	
<i>Hoplolaimus seinhorsti</i> Luc	CABI, 2012	Type 1 (CABI, 2012)	Root (CABI, 2012; Coyne et al., 2007)	No	
Mollusks					
<i>Helix aspersa</i> Muller	Zidan, 1997	Type 1 (Capinera, 2001)	Root, stem (Capinera, 2000)	No	
<i>Theba pisana</i> (Müller)	Nakhla et al., 1997	Type 1 (Garrison, 1993)	Leaves, Stem (Garrison, 1993)	No	
Viruses					
<i>Faba bean necrotic yellows nanovirus</i> (FBNYV)	Makkouk et al., 1994; Makkouk et al., 1990	Type 1 (Makkouk et al., 1990)	Leaves, stem (CABI, 2012)	No	The virus is not seed transmitted. Insect vectors are <i>Acyrtosiphon pisum</i> , <i>Aphis craccivora</i> , and <i>A. fabae</i> (Aphididae). Not transmitted by <i>Myzus persicae</i> .

Pest Name	Evidence of presence in Egypt	Host status ^a	Plant part(s) association ^b	On harvested plant part(s)? ^c	Remarks
<i>Cowpea mild mottle virus</i> (CPMMV) (Tentative Carlavirus)	Brunt et al., 1996	Type 1(Brunt et al., 1996)	Leaves, stem, possibly seed (systemic chlorosis in leaves) (CABI, 2012)	No	Seed transmission has only been shown experimentally (Brunt and Kenton, 1973). Regardless, immature seeds in green pods would be unable to transmit the virus. Vector is <i>Bemisia tabaci</i> .

^aType 1 is a natural host, i.e., a plant species that becomes infested or infected by a plant pest in nature under natural conditions (e.g., natural, cultivated and/or unmanaged plants), and the plant pest is sustained on that plant species. Type 2 is a conditional host, i.e. a plant species that is only a host or a non-host under certain conditions. Type 4 refers to situations when the plant is not a food source but serves as a fomite, which is an object or material (including a harvested plant part) that may be contaminated with a pest and that could transmit that pest from one place to another.

^bThe plant part(s) listed are those for the plant species under analysis. If the information is extrapolated, such as from plant part association on other plant species, this is noted.

^c“Yes” indicates simply that the pest has a reasonable likelihood of being associated with the harvested commodity; the level of pest prevalence on the harvested commodity (low, medium, or high) is qualitatively assessed in Risk Element A1 as part of the likelihood of introduction assessment (section 3).

2.3. Pests selected for further analysis

We identified seven arthropod pests for further analysis (Table 3); we found no pathogens or other pests that were candidates for further mitigation. All of these organisms are actionable pests for the continental United States and have a reasonable likelihood of being associated with the commodity plant part(s) at the time of harvest and remaining with the commodity, in viable form, throughout the harvesting process.

Table 3. Arthropod pests selected for further analysis.

Taxonomy	Scientific Name
Coleoptera: Bruchidae	<i>Bruchus tristis</i>
Hemiptera: Monophlebidae	<i>Icerya seychellarum</i>
Hemiptera: Pseudococcidae	<i>Maconellicoccus hirsutus</i>
Lepidoptera: Lycaenidae	<i>Lampides boeticus</i>
Lepidoptera: Noctuidae	<i>Chrysodeixis chalcites</i>
	<i>Helicoverpa armigera</i>
	<i>Spodoptera littoralis</i>

3. Assessing Pest Risk Potential

3.1. Introduction

For each pest selected for further analysis, we estimate its overall pest risk potential. Risk is described by the likelihood of an adverse event, the magnitude of the consequences, and uncertainty. In this risk assessment, we first determine for each pest if there is an endangered area within the import area. The endangered area is defined as the portion of the import area where ecological factors favor the establishment of the pest and where the presence of the pest will result in economically important losses. Once an endangered area has been determined, the overall risk of each pest is then determined by two separate components:

- 1) the likelihood of its introduction into the endangered area on the imported commodity (i.e., the likelihood of an adverse event), and
- 2) the consequences of its introduction (i.e., the magnitude of the consequences).

In general, we assess both of these components for each pest. However, if we determine that the risk of either of these components is negligible, it is not necessary to assess the other, as the overall pest risk potential would be negligible regardless of the result of the second component. In other words, if we determine that the introduction of a pest is unlikely to have unacceptable consequences, we do not assess its likelihood of being introduced. Likewise, if we determine there is negligible likelihood of a pest being introduced, we do not assess its consequences of introduction.

The likelihood and consequences of introduction are assessed using different approaches.

For the consequences of introduction, we determine if the pest meets the threshold (Yes/No) of likely causing unacceptable consequences of introduction. This determination is based on

estimating the potential consequences of introduction in terms of physical losses (rather than monetary losses). The threshold is based on a proportion of damage rather than an absolute value or amount. Pests that are like to impact at least 10 percent of the production of one or more hosts are deemed “threshold pests.”

For likelihood of introduction, which is based on the likelihoods of entry and establishment, we qualitatively assess risk using the ratings Negligible, Low, Medium, and High. The risk factors comprising the model for likelihood of introduction are interdependent and, therefore, the model is multiplicative rather than additive. Thus, if any one risk factor is rated as Negligible, then the overall likelihood will be Negligible. For the overall likelihood of introduction risk rating, we define the different categories as follows:

High: Pest introduction is highly likely to occur.

Medium: Pest introduction is possible, but for that to happen, the exact combination of required events needs to occur.

Low: Pest introduction is unlikely to occur because one or more of the required events are unlikely to happen, or the full combination of required events is unlikely to align properly in time and space.

Negligible: Pest introduction is highly unlikely to occur given the exact combination of events required for successful introduction.

3.2. Assessment results

3.2.1. *Bruchus tristis*

We determined the overall likelihood of introduction to be Negligible. We present the results of this assessment in the table below.

Determination of the portion of the continental United States endangered by *Bruchus tristis*

Climatic suitability	<i>Bruchus tristis</i> occurs in parts of Europe, southern Russia, and the Middle East (Fauna Europaea, 2012; Gentry, 1965). Based on its global distribution, <i>B. tristis</i> could establish in Plant Hardiness Zones 3 to 11 in the continental United States (Fauna Europaea, 2012; Gentry, 1965; Magarey et al., 2008).
Potential hosts at risk in PRA area	Suitable hosts for <i>B. tristis</i> grow in all of these plant hardiness zones (Gentry, 1965; Magarey et al., 2008; NASS, 2009).
Economically important hosts at risk ^a	Economically important hosts in the endangered area are beans and peas (Gentry, 1965; Magarey et al., 2008; NASS, 2009).
Pest potential on economically important hosts at risk ^a	<i>Bruchus tristis</i> is an important pest on beans and peas in some areas (Gentry, 1965). On <i>P. vulgaris</i> in Poland, seed damage from the related weevil, <i>B. rufimanus</i> , ranged from 1.2 percent to 63.5 percent depending on the cultivar (Kaniuczak, 2006). In Greece, bruchid infestation rates on legumes ranged from 2.3 percent to 57.1 percent (Bakoyannis, 1987). In Washington state, <i>B. pisorum</i> caused up to 50 percent damage on dry peas (Bragg and Burns, 2000). Based on this information, <i>B. tristis</i> could cause significant yield losses if introduced into the continental United States.
Defined Endangered Area	Bean and pea plants in plant hardiness zones 3 to 11 in the continental United States are at risk for <i>B. tristis</i> establishment (Gentry, 1965; Magarey et al., 2008; NASS, 2009).

^a As defined by ISPM No. 11, supplement 2, “economically” important hosts refers to both commercial and non-market (environmental) plants (IPPC, 2011).

Assessment of the likelihood of introduction of *Bruchus tristis* into the endangered area via the importation of fresh green beans for consumption, in pods or shelled, from Egypt

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Likelihood of Entry			
Risk Element A1: Pest prevalence on the harvested commodity (= the baseline rating for entry)	High	MU	<i>Phaseolus vulgaris</i> is a main host and <i>B. tristis</i> feeds in the seeds (Gentry, 1965; Kergoat et al., 2004). We know very little about the prevalence of this species on <i>P. vulgaris</i> in Egypt.
Risk Element A2: Likelihood of surviving post-harvest processing before shipment	High	U	Production, harvesting, and post-harvest procedures in the exporting area are not being considered as part of the assessment.
Risk Element A3: Likelihood of surviving transport and storage conditions of the consignment	High	U	We found no evidence that the standard shipping and storage conditions would lead to an increase or a decrease of the pest population.
Risk Element A: Overall risk rating for likelihood of entry	High	N/A	
Likelihood of Establishment			
Risk Element B1: Likelihood of coming into contact with host material in the endangered area	Negligible	MU	Specific information on <i>B. tristis</i> is lacking, but closely related species do not complete development before seeds have ripened. In <i>B. rufimanus</i> , “Most of the larval development and pupation occurs in the hard seeds after harvest in the store” (Medjdoub-Bensaad et al., 2007) and “The adults remain in the seeds and only emerge after sowing” (Crop Genebank Database, 2012). A similar life-history is implied for <i>B. pisorum</i> in Brindley, 1946), and <i>B. brachialis</i> “emerges from the ripe seeds” of its host plants (Bridwell, 1933). Because fresh garden beans are harvested before the seeds mature, we do not believe that fresh garden beans are a viable pathway for the introduction of <i>B. tristis</i> .
Overall Likelihood of Introduction			
Combined likelihoods of entry and establishment	Negligible	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

3.2.2. *Chrysodeixis chalcites*

We determined the overall likelihood of introduction to be Medium. We present the results of this assessment in the table below.

We determined that the establishment of *C. chalcites* in the continental United States is likely to cause unacceptable impacts. We present the results of this assessment in the table below.

Determination of the portion of the continental United States endangered by *Chrysodeixis chalcites*

Climatic suitability	This insect occurs in southern Europe, Canary Islands, Africa, Mauritius, Cape Verde Islands, and the Middle East (CABI, 2012). Based on this distribution, we estimate this insect could establish in USDA Plant Hardiness Zones 8-11.
Potential hosts at risk in PRA Area	<i>Chrysodeixis chalcites</i> is polyphagous. Numerous suitable hosts are grown throughout Plant Hardiness Zones 8-11 (see below).
Economically important hosts at risk	Economically important hosts in these plant hardiness zones include soybean, tobacco, beans, potato, cauliflower, cucumber, wheat, and corn (CABI, 2012).
Pest potential on economically important hosts at risk ^a	Quantitative data on damage is lacking. However, this species is considered as one of the most serious lepidopteran pests in many countries (CABI, 2012). Endangered and Threatened species such as <i>Solanum</i> and <i>Trifolium</i> spp. (United States Fish and Wildlife Service, 2012) may be affected in the endangered area.
Defined Endangered Area	Plant hardiness zones 8 to 11 in the continental United States are at risk for <i>C. chalcites</i> establishment

^a As defined by ISPM No. 11, supplement 2, “economically” important hosts refers to both commercial and non-market (environmental) plants (IPPC, 2011).

Assessment of the likelihood of introduction of *Chrysodeixis chalcites* into the endangered area via the importation of garden beans from Egypt

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Likelihood of Entry			
Risk Element A1: Pest prevalence on the harvested commodity (= the baseline rating for entry)	High	U	We found no information about the abundance of this insect on beans, but because it is a common and important pest in Egypt, and beans are a main host, we assume that its prevalence is high. In addition, the pest is an internal feeder and may thus not be easily detected at harvest (CABI, 2012).
Risk Element A2: Likelihood of surviving post-harvest processing before shipment	High	U	Production, harvesting, and post-harvest procedures in the exporting area are not being considered as part of the assessment.
Risk Element A3: Likelihood of surviving transport and storage conditions of the consignment	High	U	We have found no evidence that the standard shipping and storage conditions would lead to an increase or a decrease of the pest population.
Risk Element A: Overall risk rating for likelihood of entry	High	N/A	

Assessment of the likelihood of introduction of *Chrysodeixis chalcites* into the endangered area via the importation of garden beans from Egypt

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Likelihood of Establishment			
Risk Element B1: Likelihood of coming into contact with host material in the endangered area	Low	MU	Beans for consumption are usually eaten. Only a tiny fraction will be discarded, and of those, only a tiny fraction will be discarded outdoors. The life stage of the pest entering with green beans would be the larva (CABI, 2012), which has to complete its development to adulthood. Beans will start decomposing at the moment they are discarded outdoors, so pests will only have a short time to complete larval development. Then, pupal development needs to occur. After an adult female emerges, it has to find a male adult, mate successfully, and then locate a host plant. Finally, suitability of climate and availability of hosts would be limited to part of the year in most parts of the endangered area.
Risk Element B2: Likelihood of arriving in the endangered area	High	MC	More than 25 percent of the U.S. population lives in the endangered area.
Risk Element B: Combined likelihood of establishment	Medium	N/A	
Overall Likelihood of Introduction			
Combined likelihoods of entry and establishment	Medium	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

Assessment of the consequences of introduction of *Chrysodeixis chalcites* into the continental United States

Criteria	Meets criteria? (Y/N)	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Direct Impacts			
Risk Element C1: Damage potential in the endangered area	Yes	MC	<i>Chrysodeixis chalcites</i> causes “considerable” damage to tomatoes, is one of the most important noctuid pests of fodder crops in Israel, is one of the principal arthropod pests on soybean in Italy, is considered the most serious semi-looper pest attacking field fruit and vegetables in Egypt, and is a serious pest of potato in Mauritius (CABI, 2012). Consequently, <i>C. chalcites</i>

Criteria	Meets criteria? (Y/N)	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
			could cause significant economic damage in the endangered area.
Risk Element C2: Spread potential	Yes	C	<i>Chrysodeixis chalcites</i> sometimes migrates from southern to northern Europe, or to Africa (CABI, 2012). This demonstrates the ability of this species to move long distances.
Risk Element C: Pest introduction is likely to cause unacceptable direct impacts	Yes	N/A	
Conclusion			
Is the pest likely to cause unacceptable consequences in the PRA area?	Yes	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

3.2.3. *Helicoverpa armigera*

We determined the overall likelihood of introduction to be Medium. We present the results of this assessment in the table below.

We determined that the establishment of *H. armigera* in the continental United States is likely to cause unacceptable impacts. We present the results of this assessment in the table below.

Determination of the portion of the continental United States endangered by *Helicoverpa armigera*

Climatic suitability	This insect is widely distributed and occurs throughout Europe, the Middle East, Central and South Asia, the Far East, Africa, Australia, and Oceania (CABI, 2012). Based on this distribution, we estimate establishment is possible in USDA Plant Hardiness Zones 5-11.
Potential hosts at risk in PRA Area	Numerous suitable hosts are grown throughout plant hardiness zones 5-11 (see below).
Economically important hosts at risk	Economically important hosts in plant hardiness zones 5-11 include cotton, pigeon pea, chickpea, tomato, okra, pea, soybean, tobacco, potato, corn, and citrus (CABI, 2012).
Pest potential on economically important hosts at risk ^a	<i>Helicoverpa armigera</i> is a serious economic pest. In India, it routinely destroys over half of the pigeon pea and chickpea crops (Reed and Pawar, 1982), and 10-100 percent damage has been reported on potato (Parihar and Singh, 1988). In New Zealand, an outbreak of this pest once caused major damage to <i>Pinus radiata</i> trees (50 percent of foliage consumed on over 60 percent of the trees) (CABI, 2012). The pest is also economically very important in several European countries (CABI, 2012).

Defined Endangered Area	Cotton, tomato, okra, pea, soybean, tobacco, potato, corn, citrus, and other crops (see above) in plant hardiness zones 5 to 11 in the continental United States are at risk for <i>H. armigera</i> establishment.
--------------------------------	--

^a As defined by ISPM No. 11, supplement 2, “economically” important hosts refers to both commercial and non-market (environmental) plants (IPPC, 2011).

Assessment of the likelihood of introduction of *Helicoverpa armigera* into the endangered area via the importation of garden beans from Egypt

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Likelihood of Entry			
Risk Element A1: Pest prevalence on the harvested commodity (= the baseline rating for entry)	High	MC	<i>Helicoverpa armigera</i> is widespread in Egypt and <i>P. vulgaris</i> is among the main hosts of this pest (CABI, 2012).
Risk Element A2: Likelihood of surviving post-harvest processing before shipment	High	U	Production, harvesting, and post-harvest procedures in the exporting area are not being considered as part of the assessment.
Risk Element A3: Likelihood of surviving transport and storage conditions of the consignment	High	U	We found no evidence that the standard shipping and storage conditions would lead to an increase or a decrease of the pest population.
Risk Element A: Overall risk rating for likelihood of entry	High	N/A	
Likelihood of Establishment			
Risk Element B1: Likelihood of coming into contact with host material in the endangered area	Low	MC	Beans for consumption are usually eaten. Only a tiny fraction will be discarded, and of those, only a tiny fraction will be discarded outdoors. The life stage of the pest entering with green pods would be larvae (CABI, 2012), which have to complete development before they could fly to a host plant. Beans will start decomposing as soon as they are discarded, so larval development would have to be completed in a short time. Then, pupal development needs to successfully occur. After a female emerges, it would have to find an adult male, mate successfully, and then find a host plant. Finally, suitability of climate and availability of hosts would be limited to part of the year in most parts of the endangered area.
Risk Element B2: Likelihood of arriving in the endangered area	High	MC	More than 25 percent of the U.S. population lives in the endangered area.
Risk Element B: Combined	Medium	N/A	

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
likelihood of establishment			
Overall Likelihood of Introduction			
Combined likelihoods of entry and establishment	Medium	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

Assessment of the consequences of introduction of *Helicoverpa armigera* into the continental United States

Criteria	Meets criteria? (Y/N)	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Direct Impacts			
Risk Element C1: Damage potential in the endangered area	Yes	MC	As discussed above (Endangered Area table), this pest can be a serious economic pest.
Risk Element C2: Spread potential	Yes	C	In South Africa, <i>H. armigera</i> lays an average of 730 eggs over an oviposition period of 10-23 days; females may produce up to 11 generations per year in tropical climates. Larvae have limited mobility, but adults can fly long distances (CABI, 2012).
Risk Element C: Pest introduction is likely to cause unacceptable direct impacts	Yes	N/A	
Conclusion			
Is the pest likely to cause unacceptable consequences in the PRA area?	Yes	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

3.2.4. *Icerya seychellarum*

We determined the overall likelihood of introduction to be Negligible. We present the results of this assessment in the table below.

Determination of the portion of the PRA Area endangered by *Icerya seychellarum*

Climatic suitability	This pest is distributed throughout much of subtropical and tropical Asia. It also occurs in several African countries, Oceania, and some countries in South America (CABI, 2012). Based on its global distribution, this pest could survive in U.S. Plant Hardiness Zones 9-11.
Potential hosts at risk in PRA Area	The pest feeds on a large number of hosts in various families (CABI, 2012) (see below). Many of its hosts are widely distributed and abundant in the continental United States, including plant hardiness zones 9-11.
Economically important hosts at risk	Economically important hosts growing in plant hardiness zones 9-11 in the United States include peach, pomegranate, mango, guava, avocado, fig, date,

	paw paw, grape, sweet potato, tomato, lettuce, and beans (CABI, 2012).
Pest potential on economically important hosts at risk ^a	<i>Icerya seychellarum</i> killed fruit trees in the Pacific Islands (Williams and Watson, 1990) and is a pest of fruit trees in the Seychelles, the Mascarene Islands, and Japan (CABI, 2012).
Defined Endangered Area	Peach, pomegranate, avocado, grape, sweet potato, tomato, lettuce, and bean crops in plant hardiness zones 9 to 11 in the continental United States are at risk.

^a As defined by ISPM No. 11, supplement 2, “economically” important hosts refers to both commercial and non-market (environmental) plants (IPPC, 2011).

Assessment of the likelihood of introduction of *Icerya seychellarum* into the endangered area via the importation of garden beans from Egypt

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Likelihood of Entry			
Risk Element A1: Pest prevalence on the harvested commodity (= the baseline rating for entry)	Low	MU	We found no information about the prevalence of <i>I. seychellarum</i> on beans. In other crops, pest populations seem to be generally low (Newberry and Hill, 1985), perhaps because of natural enemies (Hill, 1983). Therefore, we rated the prevalence of <i>I. seychellarum</i> as low.
Risk Element A2: Likelihood of surviving post-harvest processing before shipment	Low	U	For this risk assessment, we assume that no post-harvest processing takes place. We also have no evidence that the population could increase during the time between harvest and transport.
Risk Element A3: Likelihood of surviving transport and storage conditions of the consignment	Low	U	We found no evidence that the standard shipping and storage conditions would lead to an increase or a decrease of the pest population.
Risk Element A: Overall risk rating for likelihood of entry	Low	N/A	
Likelihood of Establishment			
Risk Element B1: Likelihood of coming into contact with host material in the endangered area	Negligible	MC	Beans for consumption are usually eaten. Only a tiny fraction will be discarded, and of those, only a tiny fraction will be discarded outdoors. Adult <i>I. seychellarum</i> are immobile; only the immature stage (crawler) can move on its own, but only over very short distances (generally less than 1m) (Magsig-Castillo et al., 2010). Crawlers can disperse over long distances by wind, but that requires climbing high enough to encounter suitable air movement, and then being blown onto a

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
			suitable host plant (CABI, 2012). Additionally, crawler production would have to coincide with when the pod is discarded. Finally, suitability of climate and availability of hosts are limited to parts of the year in most of the endangered area. Overall, because the pest life stage present is highly unlikely to develop into a dispersing life stage, and that life stage is highly unlikely to move on its own from the commodity to a new host, we rated this element Negligible.
Risk Element B2: Likelihood of arriving in the endangered area	N/A	N/A	
Risk Element B: Combined likelihood of establishment	Negligible	N/A	
Overall Likelihood of Introduction			
Combined likelihoods of entry and establishment	Negligible	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

3.2.5. *Lampides boeticus*

We determined the overall likelihood of introduction to be Medium. We present the results of this assessment in the table below.

We determined that the establishment of *L. boeticus* in the continental United States is likely to cause unacceptable impacts. We present the results of this assessment in the table below.

Determination of the portion of the continental United States endangered by *Lampides boeticus*

Climatic suitability	This pest is present in southern Europe, Africa, Oceania, the Middle East, and southern Asia (CABI, 2012). Based on this distribution, we estimate it could establish in USDA Plant Hardiness Zones 8-11.
Potential hosts at risk in PRA Area	Suitable hosts for <i>L. boeticus</i> , including various types of beans and peas (CABI, 2012), are grown in all of these plant hardiness zones.
Economically important hosts at risk	Economically important hosts in these plant hardiness zones include soybean, snap bean, lima bean, sweet pea, and Lucerne (NASS, 2009; CABI, 2012).
Pest potential on economically important hosts at risk ^a	In Hawaii, <i>L. boeticus</i> is a major pest of garden beans (CABI, 2012). In India, damage to pods and locules of peas averaged 8 percent, which was considered significant (CABI, 2012).
Defined Endangered Area	Soybean, snap bean, lima bean, sweet pea, and lucerne crops in plant hardiness zones 8 to 11 in the continental United States are at risk.

^a As defined by ISPM No. 11, supplement 2, "economically" important hosts refers to both commercial and non-market (environmental) plants (IPPC, 2011).

Assessment of the likelihood of introduction of *Lampides boeticus* into the endangered area via the importation of garden beans from Egypt

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Likelihood of Entry			
Risk Element A1: Pest prevalence on the harvested commodity (= the baseline rating for entry)	High	MC	<i>Lampides boeticus</i> is common in Egypt, where it was one of the most abundant insects in alfalfa fields (Shebl et al., 2009) and a dominant species in cowpeas (Abdel-Rahman and Amro, 2004). <i>Phaseolus vulgaris</i> is among the preferred hosts of this pest.
Risk Element A2: Likelihood of surviving post-harvest processing before shipment	High	U	Production, harvesting, and post-harvest procedures in the exporting area are not being considered as part of the assessment.
Risk Element A3: Likelihood of surviving transport and storage conditions of the consignment	High	U	We found no evidence that the standard shipping and storage conditions would lead to an increase or a decrease of the pest population.
Risk Element A: Overall risk rating for likelihood of entry	High	N/A	
Likelihood of Establishment			
Risk Element B1: Likelihood of coming into contact with host material in the endangered area	Low	MC	Beans for consumption are usually eaten. Only a tiny fraction will be discarded, and of those, only a tiny fraction will be discarded outdoors. The life stage entering with green pods would be larvae (CABI, 2012), which have to complete development before they could fly to a host plant. Beans will start decomposing as soon as they are discarded, so larval development would have to be completed in a short time. Then, pupal development needs to successfully occur. After a female emerges, it would have to find an adult male, mate successfully, and then find a host plant. Finally, suitability of climate and availability of hosts would be limited to part of the year in most parts of the endangered area.
Risk Element B2: Likelihood of arriving in the endangered area	High	MC	More than 25 percent of the U.S. population lives in the endangered area.
Risk Element B: Combined likelihood of establishment	Medium	N/A	
Overall Likelihood of Introduction			
Combined likelihoods of entry and establishment	Medium	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

Assessment of the consequences of introduction of *Lampides boeticus* into the continental United States

Criteria	Meets criteria? (Y/N)	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Direct Impacts			
Risk Element C1: Damage potential in the endangered area	Yes	MU	<i>Lampides boeticus</i> is a major pest of green beans in Hawaii (CABI, 2012). Green beans are grown throughout all of the endangered area.
Risk Element C2: Spread potential	Yes	C	<i>Lampides boeticus</i> can fly long distances (CABI, 2012), and routinely migrates from southern to northern Europe, and to Africa. Moreover, the species has established in New Zealand and Hawaii (CABI, 2012).
Risk Element C: Pest introduction is likely to cause unacceptable direct impacts	Yes	N/A	
Conclusion			
Is the pest likely to cause unacceptable consequences in the PRA area?	Yes	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

3.2.6. *Maconellicoccus hirsutus*

We determined the overall likelihood of introduction to be Negligible. We present the results of this assessment in the table below.

Determination of the portion of the PRA Area endangered by *Maconellicoccus hirsutus*

Climatic suitability	<i>Maconellicoccus hirsutus</i> is present in many parts of Asia, Africa, Oceania, the Middle East, and the Caribbean and in some parts of the southern and western United States (CABI, 2012). Based on this distribution, we estimate that it may establish in the United States throughout Plant Hardiness Zones 7-11.
Potential hosts at risk in PRA Area	The pest feeds on a large number of hosts in various families, including ornamentals and native plants (CABI, 2012). Many of its hosts are widely distributed and abundant in the continental United States.
Economically important hosts at risk	Numerous economically important hosts grow in plant hardiness zones 7-11 in the United States, including citrus, avocado, cherry, plum, pepper, grapes, corn, beans and peas, eggplant, potato, cucumber, cabbage, squash, okra, and tomato (CABI, 2012).
Pest potential on economically important hosts at risk ^a	In India, <i>M. hirsutus</i> has caused economic losses in cotton, grapevine, mulberry, and pigeonpea (CABI, 2012). These hosts occur in the United States in Plant Hardiness Zones 7-11.

Defined Endangered Area	The crops mentioned above are at risk in plant hardiness zones 7 to 11 in the continental United States, except for those states where the pest has already established (California, Florida, Texas) (Horton, 2008; National Invasive Species Council, No Date; Stang, 2009).
--------------------------------	---

^a As defined by ISPM No. 11, supplement 2, “economically” important hosts refers to both commercial and non-market (environmental) plants (IPPC, 2011).

Assessment of the likelihood of introduction of *Maconellicoccus hirsutus* into the endangered area via the importation of garden beans from Egypt

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Likelihood of Entry			
Risk Element A1: Pest prevalence on the harvested commodity (= the baseline rating for entry)	Low	MC	<i>Maconellicoccus hirsutus</i> is likely to be highly visible on green beans because the pest forms colonies that become covered by white, woolly, wax material (CABI, 2012). Routinely harvesting and exporting even moderately infested beans would not be commercially viable and is unlikely to occur. Therefore, pests are only likely to occur on harvested beans at low (less visible) densities.
Risk Element A2: Likelihood of surviving post-harvest processing before shipment	Low	U	For this risk assessment, we assume that no post-harvest processing takes place. We also have no evidence that the population could increase during the time between harvest and transport.
Risk Element A3: Likelihood of surviving transport and storage conditions of the consignment	Low	U	We found no evidence that the standard shipping and storage conditions would lead to an increase or a decrease of the pest population.
Risk Element A: Overall risk rating for likelihood of entry	Low	N/A	
Likelihood of Establishment			
Risk Element B1: Likelihood of coming into contact with host material in the endangered area	Negligible	MC	Beans for consumption are usually eaten. Only a tiny fraction will be discarded, and of those, only a tiny fraction will be discarded outdoors. Adult <i>M. hirsutus</i> are immobile; only the immature stage (crawler) can disperse, but only over very short distances (Caribbean Pest Information Network, No Date). Crawlers can disperse over long distances by wind, but that requires climbing high enough to encounter suitable air movement, and then being blown onto a suitable host plant (CABI, 2012). Additionally, crawler production would

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
			have to coincide with when the pod is discarded. Finally, suitability of climate and availability of hosts are limited to parts of the year in most of the endangered area. Overall, because the pest life stage present is highly unlikely to develop into a dispersing life stage, and that life stage is highly unlikely to move on its own from the commodity to a new host, we rated this element Negligible.
Risk Element B2: Likelihood of arriving in the endangered area	N/A	N/A	
Risk Element B: Combined likelihood of establishment	Negligible	N/A	
Overall Likelihood of Introduction			
Combined likelihoods of entry and establishment	Negligible	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

3.2.7. *Spodoptera littoralis*

We determined the overall likelihood of introduction to be Medium. We present the results of this assessment in the table below.

We determined that the establishment of *S. littoralis* in the continental United States is likely to cause unacceptable impacts. We present the results of this assessment in the table below.

Determination of the portion of the continental United States endangered by *Spodoptera littoralis*

Climatic suitability	The insect is recorded in Africa, southern Europe, and the Middle East (CABI, 2012). Therefore, we estimate it could establish in USDA Plant Hardiness Zones 8-11.
Potential hosts at risk in PRA Area	The host range of <i>S. littoralis</i> comprises over 40 families, and suitable hosts are present throughout plant hardiness zone 8-11.
Economically important hosts at risk	Economically important hosts in these plant hardiness zones include cotton, tobacco, potato, tomato, onion, citrus, beans, peppers, grapes, alfalfa, and various grasses (CABI, 2012).
Pest potential on economically important hosts at risk ^a	EPPO (2006a) states the following: <ul style="list-style-type: none"> • <i>Spodoptera littoralis</i> is one of the most destructive agricultural lepidopterous pests within its subtropical and tropical range. • It is polyphagous and attacks a number of economically important hosts. • On cotton, the pest may cause considerable damage. • Pods and seeds of cowpeas are often badly damaged. • In tomatoes, larvae bore into the fruit, making them unmarketable.

	<ul style="list-style-type: none"> • In Italy, it is an important pest of protected crops of ornamentals and vegetables.
Defined Endangered Area	Cotton, tobacco, potato, tomato, onion, citrus, beans, peppers, grapes, alfalfa, and various grass crops in plant hardiness zones 8 to 11 in the continental United States are at risk for <i>S. littoralis</i> establishment.

^a As defined by ISPM No. 11, supplement 2, “economically” important hosts refers to both commercial and non-market (environmental) plants (IPPC, 2011).

Assessment of the likelihood of introduction of *Spodoptera littoralis* into the endangered area via the importation of garden beans from Egypt

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Likelihood of Entry			
Risk Element A1: Pest prevalence on the harvested commodity (= the baseline rating for entry)	High	MC	<i>Spodoptera littoralis</i> is widespread in Egypt (CABI, 2012). <i>Phaseolus vulgaris</i> is among the preferred hosts of this pest.
Risk Element A2: Likelihood of surviving post-harvest processing before shipment	High	U	Production, harvesting, and post-harvest procedures in the exporting area are not being considered as part of the assessment.
Risk Element A3: Likelihood of surviving transport and storage conditions of the consignment	High	U	We found no evidence that the standard shipping and storage conditions would lead to an increase or a decrease of the pest population.
Risk Element A: Overall risk rating for likelihood of entry	High	N/A	
Likelihood of Establishment			
Risk Element B1: Likelihood of coming into contact with host material in the endangered area	Low	MC	Beans for consumption are usually eaten. Only a tiny fraction will be discarded, and of those, only a tiny fraction will be discarded outdoors. The life stage entering with green pods would be larvae (CABI, 2012), which have to complete development before they could fly to a host plant. Beans will start decomposing as soon as they are discarded, so larval development would have to be completed in a short time. Then, pupal development needs to successfully occur. After a female emerges, it would have to find an adult male, mate successfully, and then find a host plant. Finally, suitability of climate and availability of hosts would be limited to part of the year in most parts of the endangered area.
Risk Element B2: Likelihood of arriving in the endangered area	High	MC	More than 25 percent of the U.S. population lives in the endangered area.

Risk Element	Risk Rating	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Risk Element B: Combined likelihood of establishment	Medium	N/A	
Overall Likelihood of Introduction			
Combined likelihoods of entry and establishment	Medium	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

Assessment of the consequences of introduction of *Spodoptera littoralis* into the continental United States

Criteria	Meets criteria? (Y/N)	Uncertainty Rating ^a	Justification for rating and explanation of uncertainty (and other notes as necessary)
Direct Impacts			
Risk Element C1: Damage potential in the endangered area	Yes	MC	As discussed above (Endangered Area table), this pest seems likely to cause economic damage in the continental United States if introduced, despite a lack of more specific, quantitative information.
Risk Element C2: Spread potential	Yes	C	The flight range of this pest can be up to 1.5 km in four hours (EPPO, 2006a). In optimal climates, the pest can have up to 7 overlapping generations per year, with an average of 20-1000 eggs produced by each female (CABI, 2012).
Risk Element C: Pest introduction is likely to cause unacceptable direct impacts	Yes	N/A	
Conclusion			
Is the pest likely to cause unacceptable consequences in the PRA area?	Yes	N/A	

^a C=Certain, MC=Moderately Certain, MU=Moderately Uncertain, U=Uncertain

4. Summary and Conclusions of Risk Assessment

Of the organisms associated with green beans worldwide and reported in Egypt, we identified organisms that are actionable pests for the continental United States and have a reasonable likelihood of being associated with the commodity following harvesting from the field and prior to any post-harvest processing. We evaluated these organisms for their likelihood of introduction (i.e., entry plus establishment) and their potential consequences of introduction. Pests that meet the threshold to likely cause unacceptable consequences of introduction and receive an overall likelihood of introduction risk rating above Negligible are candidates for risk management. The results of this risk assessment represent a baseline estimate of the risks associated with the

import commodity pathway as described in section 1.4. (i.e., green beans in pods or shelled with no mandatory production or post-harvest processes).

Of the pests selected for further analysis, we determined that those identified below (Table 4) are *not* candidates for risk management, either because no portion of the continental United States is likely to be endangered by the pest, they do not meet the threshold to likely cause unacceptable consequences of introduction, and/or because they received a Negligible overall risk rating for likelihood of introduction into the endangered area via the import pathway. We summarize the results for each pest below (Table 4).

All the other pests selected for further analysis are candidates for risk management, because they meet the threshold to likely cause unacceptable consequences of introduction, and they received an overall likelihood of introduction risk rating above Negligible. We summarize the results for each pest below (Table 5).

Detailed examination and choice of appropriate phytosanitary measures to mitigate pest risk are part of the pest risk management phase within APHIS and are not addressed in this document.

Table 4. Summary for pests selected for further evaluation and determined *not* to be candidates for risk management

Pest	Endangered area within the PRA area	Meets unacceptable Consequences of Introduction threshold	Likelihood of Introduction overall rating
<i>Bruchus tristis</i>	Yes	N/A	Negligible
<i>Icerya seychellarum</i>	Yes	N/A	Negligible
<i>Maconellicoccus hirsutus</i>	Yes	N/A	Negligible

Table 5. Summary for pests selected for further evaluation and determined to be candidates for risk management (All meet the threshold for unacceptable consequences of introduction.)

Pest	Likelihood of Introduction overall rating
<i>Lampides boeticus</i>	Medium
<i>Chrysodeixis chalcides</i>	Medium
<i>Helicoverpa armigera</i>	Medium
<i>Spodoptera littoralis</i>	Medium

5. Acknowledgements

Authors Heike Meissner, Risk Analyst^a
Feridoon Mehdizadegan, Risk Analyst^a
Glenn Fowler, Risk Analyst^a

Reviewers Alison Neeley, Risk Analyst^a
John Rogers, Risk Analyst^a

^a Plant Epidemiology and Risk Analysis Laboratory, USDA-APHIS-PPQ

6. Literature Cited

- 7 CFR § 319.56. 2012. U.S. Code of Federal Regulations, Title 7, Part 319 (7 CFR § 319.56 - Fruits and Vegetables).
- Abate, T., and J. K. O. Ampofo. 1996. Insect pests of beans in Africa: their ecology and management. *Annual Review of Entomology* 41:45-73.
- Abd-Rabou, S. 2006. Importation, colonization and establishment of the parasitoid, *Encarsia pergandiella* Howard (Hymenoptera: Aphelinidae) on *Bemisia argentifolii* Bellows & Perring (Homoptera: Aleyrodidae) in Egypt. *Journal of Biological Control* 20(2):119-122.
- Abd-Rabou, S., and N. Ahmed. 2008. Bionomics of *Bemisia afer* (Hemiptera: Aleyrodidae) A New Pest of *Citrus aurantium* var. *amara* in Egypt [Abstract]. *Egyptian Journal of Agricultural Research* 86(5):1783.
- Abdel-Dali, M. 2007. "*Phseolus vulgaris* Pests". Personal communication to Nancy Arciniegas (CPHST Colombia Pest Risk Analyst) on August 21, 2007, from Dr. Marwan Abdel Dali NCARTT (National Center for Agricultural Research and Technology Transfer).
- Abdel-Moniem, A. S. H., T. E. A. El-Wahab, and N. A. Farag. 2011. Prevailing insects in Roselle plants, *Hibiscus sabdariffa* L., and their efficiency on pollination [ABSTRACT]. *Archives of Phytopathology and Plant Protection* 44(3):242-252.
- Abdel-Rahman, M., and M. Amro. 2004. Incidence of certain arthropod pests and predators inhabiting cowpea, with special reference to the varietal resistance of selected cultivars to *Bemisia tabaci* (Gen.) and *Tetranychus urticae* Koch Ass. Univ. Bull. Environ. Res. 7(1):31-39.
- Abdel Rahman, K. M. 2001. Food consumption and utilisation of the grasshopper *Chrotogonus lugubris* Blanchard (Orthoptera, Acridoidea, Pyrgomorphidae) and its effect on the egg deposition. *Journal of Central European Agriculture* 2(No. 3-4):263-270.
- Afifi, F. M. L., and H. M. A. El-Whab. 1990. The biology of the cotton leaf-worm, *Spodoptera littoralis* (Boisd.), (Lepidoptera: Noctuidae), on three leguminous summer vegetable crops. *Bulletin of Faculty of Agriculture, University of Cairo* 41(3):609-620.
- Al-Azawi, A. F. 1967. Agromyzid leafminers and their parasites in Iraq. *Bulletin of Entomological Research* 57(2):285-287.
- Allen, C. 2012. *Ralstonia solanacearum* Personal communication to F. Mehdizadegan on August, 20, 2012, from
- Álvarez, B., J. Vasse, V. Le-Courtois, D. Trigalet-Démery, M. M. López, and A. Trigalet. 2007. Comparative Behavior of *Ralstonia solanacearum* Biovar 2 in Diverse Plant Species. *Phytopathology* 98(1):59-68.
- Ammar, E. D., S. Elnagar, A. E. Abul-Ata, and G. H. Sewify. 1989. Vector and host-plant relationships of the leafhopper-borne maize yellow stripe virus. *Journal of Phytopathology* 126(3):246-252.
- Ampofo, K. 1994. Field Pests of Bean in Africa. *Centra Internacional de Agricultura Tropical*, Cali, Colombia.
- Anonymous. 1978. Report of the Crop Protection Services, Department of Agriculture, Gambia [ABSTRACT] (Annual report 1977). Gambia, Department of Agriculture, Crop Protection Services. 40 pp.
- APHIS. 2012. Fruits and Vegetables Import Requirements (FAVIR). United States Department of Agriculture, Animal and Plant Health Inspection Service (APHIS). <https://epermits.aphis.usda.gov/manual/index.cfm?action=pubHome>. (Archived at PERAL).

- Arnett, R. H. J., M. C. Thomas, P. E. Skelley, and J. H. Frank. 2002. American Beetles Volume 2: Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press LCC, Boca Raton, Florida. 861 pp.
- Ascher, K. R. S., M. Streloke, G. H. Schmidt, and J. D. Warthen Jr. 1989. The antifeedant effect on neem seed kernel extract and azadirachtin on nymphs of *Eyprepocnemis plorans* [ABSTRACT]. *Phytoparasitica* 17:167-174.
- Avidov, Z., and I. Harpaz. 1969. Plant pests of Israel. Israel Universities Press, Jerusalem. 549 pp.
- Bakoyannis, A. E. 1987. Bruchids damaging leguminous plants in the prefectures of Magnesia and Larissa. *Georgike Ereuna* 11(4):429-435.
- Bellinger, P. F., K. A. Christiansen, and G. Janssens. 2012. Checklist of the Collembola of the World. <http://www.collembola.org/>. (Archived at PERAL).
- Ben-Dov, Y., D. R. Miller, and G. A. P. Gibson. 2012. ScaleNet. <http://www.sel.barc.usda.gov/scalenet/scalenet.htm>. (Archived at PERAL).
- Bisby, F., Y. Roskov, A. Culham, T. Orrell, D. Nicolson, L. Paglinawan, N. Bailly, W. Appeltans, P. Kirk, T. Bourgoin, G. Baillargeon, and D. Ouvrard (eds.). 2012. Species 2000 & ITIS Catalogue of Life, 2012 Annual Checklist. Species 2000: Reading, UK. www.catalogueoflife.org/col/. (Archived at PERAL).
- Bradbury, J. F. 1986. Guide to Plant Pathogenic Bacteria. CAB International Mycological Institute. Ferry Lane, Kew, Surrey, England.
- Bragg, D., and G. W. Burns. 2000. Crop profile for peas (dry) in Washington. WSU Cooperative Extension Tri-County Area; Crop and Soil Sciences Department Washington State University. 10 pp.
- Bridwell, J. C. 1933. The hairy-vetch bruchid, *Bruchus brachialis* Fahrhaeus, in the United States. *Journal of Agricultural Research* 46(8):739-751.
- Brindley, T. A. 1946. Pea weevil. *Farmers' Bulletin* 1971.
- Brown, J. K. 2007. The *Bemisia tabaci* Complex: Genetic and Phenotypic Variability Drives Begomovirus Spread and Virus Diversification. American Phytopathological Society. 18 pp.
- Brunt, A. A., K. Crabtree, M. J. Dallwitz, A. J. Gibbs, L. Watson, and E. J. Zurcher. 1996. Plant Viruses Online: Descriptions and Lists from the VIDE Database. Version: 20th <http://biology.anu.edu.au/Groups/MES/vide/>. (Archived at PERAL).
- Brunt, A. A., and R. Kenton. 1973. Cowpea mild mosaic virus, a newly recognized virus infecting cowpea (*Vigna unguiculata*) in Ghana. *Ann. Appl. Biol.* 74:67-74.
- C.A.B. International. 1987. Distribution Maps of Plant Pests: *Agrotis segetum* (Denis & Schiffermueller) (Map no. 490). C.A.B. International, London. 3 pp.
- CABI. 2012. Crop Protection Compendium. Commonwealth Agricultural Bureau International (CABI). <http://www.cabi.org/cpc/>. (Archived at PERAL).
- CABI/EPPO. 1997. Distribution Maps of Plant Pests: *Maconellicoccus hirsutus* (Map No. 100 2nd Revision). CAB International (CABI), European and Mediterranean Plant Protection Organization (EPPO), Wallingford, UK, and Paris, France. 4 pp.
- CABI/EPPO. 2008. Distribution Maps of Plant Pests: *Icerya seychellarum* (Map No. 52 1st Revision). CAB International (CABI), European and Mediterranean Plant Protection Organization (EPPO), Wallingford, UK, and Paris, France. 4 pp.
- Canerday, T. D., and F. S. Arant. 1964. The effect of late season infestations of the strawberry spider mite, *Tetranychus atlanticus*, on cotton production [ABSTRACT]. *Journal of Economic Entomology* 57(6):931-933.
- Capinera, J. L. 2000. *Ostrinia nubilalis* (Hübner) (Insecta: Lepidoptera: Pyralidae) EENY-156. Florida Department of Agriculture and Consumer Services, Division of Plant Industry. University of Florida. Last accessed January 10, 2008, <http://pick4.pick.uga.edu/mp/20q?search=Ostrinia+nubilalis>.
- Capinera, J. L. 2001. Handbook of Vegetable Pests. Academic Press, San Diego. 729 pp.

- Caribbean Pest Information Network. No Date. Pests of Phytosanitary Significance to the Caribbean Sub-Region: *Maconellicoccus hirsutus*. http://www.caripestnetwork.org/vtt/docs/datasheets/hemiptera_homoptera/maconellicoccus_hirsutus.pdf. (Archived at PERAL).
- Childers, C. C., J. C. V. Rodrigues, and W. C. Welbourn. 2003. Host plants of *Brevipalpus californicus*, *B. obovatus*, and *B. phoenicis* (Acari: Tenuipalpidae) and their potential involvement in the spread of viral diseases vectored by these mites. *Experimental and Applied Acarology* 30(1-3):29-105.
- Commonwealth Institute of Entomology. 1977. Distribution Maps of Plant Pests: *Chrysodeixis chalcites* (Esp.) (Map No. 376). Commonwealth Agricultural Bureaux, London. 2 pp.
- Commonwealth Institute of Entomology. 1984. Distribution Maps of Plant Pests: *Lampides boeticus* (Linnaeus) (Map no. 459). Commonwealth Agricultural Bureaux, London. 3 pp.
- Coyne, D. L., J. M. Nicol, and B. Claudius-Cole. 2007. Practical plant nematology: a field and laboratory guide.:93. Last accessed August 3, 2012, http://www.spipm.cgiar.org/c/document_library/get_file?p_l_id=17829&folderId=18466&name=DLFE-138.pdf.
- Creamer, R., M. Luque-Williams, and M. Howo. 1996. Epidemiology and incidence of Beet Curly Top Gemenivirus in naturally infected weed hosts. *Plant dis.* 80:533-535.
- Crop Genebank Database. 2012. Bean seed beetle. http://cropgenebank.sgrp.cgiar.org/index.php?option=com_content&view=article&id=557%3Ainsects-for-faba-bean&catid=166%3Astog-faba-bean&Itemid=752&lang=es. (Archived at PERAL).
- De Meyer, M., S. Mohamed, and I. M. White. 2012. Invasive Fruit Fly Pests in Africa. Last accessed October 24, 2012, <http://www.africamuseum.be/fruitfly/AfroAsia.htm>.
- Duffy, S., and E. Holmes. 2007. Multiple Introductions of the Old World Begomovirus Tomato yellow leaf curl virus into the New World. *Appl Environ Microbiol* 21:7114-7117.
- EcoPort. 2012. *Schistocerca gregaria* EcoPort Foundation. <http://ecoport.org>. (Archived at PERAL).
- El-Dessouki, S. A., and M. M. Hosny. 1969. Host-plants, symptoms of infestation and certain characteristics of *Empoasca* spp. (Jassidae) in the Cairo area, U.A.R. *Zeitschrift fur Angewandte Entomologie* 63(pt. 3):272-280.
- El-Lakwa, F. A. M., E. F. El-Khayat, A. A. Hafez, and H. H. Shalaby. 1999. Susceptibility of three varieties of bean (*Phaseolus vulgaris* L.) to infestation with whitefly and aphids. *Annals of Agricultural Science, Moshtohor* 37(1):585-603.
- El-Minshawy, A. M., and N. H. El-Hinnawy. 1976. Population fluctuations of grasshoppers and locusts in Alexandria area (Egypt) (Orthoptera: Acridiidae [sic] and Tettigoniidae) [ABSTRACT]. *Bulletin de la Société Entomologique d'Egypte* 59:83-98.
- El Kifl, A. H., A. E. A. Wahab, M. A. Assem, and A. A. Metwally. 1974. List of insects, mites and pests associated with leguminous crops in Egypt. *Bull. Soc. Ent. Egypte* 58:297-302.
- Elbadry, B. E., G. M. Mousa, and E. M. Bakr. 2006. Pesticidal efficiency of newly synthesized organo-cyanide compounds against certain pests infested bean plants. *Egyptian Journal of Agricultural Research* 84(1):101-110.
- EPPO. 2000. Reporting Service 2000 (No. 4). European and Mediterranean Plant Protection Organization (EPPO), Paris. 26 pp.
- EPPO. 2004. EPPO Standards: Diagnostic protocols for regulated pests. *Bemisia tabaci* (PM 7/35(1)). European and Mediterranean Plant Protection Organization (EPPO), Paris. *EPPO Bulletin* 34(2):281-288.
- EPPO. 2005. Plant quarantine information retrieval system. Ver 4.5. European and Mediterranean Plant Protection Organization (EPPO), Paris.
- EPPO. 2006a. Datasheets on quarantine pests: *Spodoptera littoralis* and *Spodoptera litura*. European and Mediterranean Plant Protection Organization (EPPO), Paris.

- EPPO. 2006b. Distribution Maps of Quarantine Pests for Europe: *Liriomyza bryoniae*. European and Mediterranean Plant Protection Organization (EPPO), Paris. 2 pp.
- EPPO. 2009. EPPO Reporting Service. European and Mediterranean Plant Protection Organization (EPPO), Paris.
- EPPO. 2012a. EPPO Global Database. European and Mediterranean Plant Protection Organization (EPPO), Paris. (Archived at PERAL).
- EPPO. 2012b. Datasheets on Quarantine Pests: Tomato yellow leaf curl bigeminivirus. European and Mediterranean Plant Protection Organization (EPPO), Paris.
http://www.eppo.int/QUARANTINE/virus/TYLC_virus/TYLCV0_ds.pdf. (Archived at PERAL).
- Farr, D. F., A. Y. Rossman, M. E. Palm, and E. B. McCray. 2007. Fungal Databases. United States Department of Agriculture, Agricultural Research Service, Systematic Mycology and Microbiology Laboratory. <http://nt.ars-grin.gov/fungaldatabases/>. (Archived at PERAL).
- Farrag, R. M., and E. A. Zakzouk. 2000. Relative abundance of some pests and their control on some host plants. *Egyptian Journal of Agricultural Research* 78(5):1897-1904.
- Fauna Europaea. 2012. *Bruchus tristis* Bohemann. http://www.faunaeur.org/distribution_table.php. (Archived at PERAL).
- Favret, C. 2012. Aphid Species File. Version 1.0/4.1. <http://Aphid.SpeciesFile.org>. (Archived at PERAL).
- Ferris, H. 2012. The Nematode-Plant Expert Information System NEMAPLEX. The Virtual Encyclopedia on Soil and Plant Nematodes. University of California, Department of Nematology.
<http://plpnemweb.ucdavis.edu/nemaplex/Taxadata/g138s9.htm>. (Archived at PERAL).
- Garrison, R. W. 1993. New Agricultural Pest for Southern California White Garden Snail (*Theba pisana*). LA County Agricultural Commissioner/Weights and Measures Department. Last accessed March 25, 2007, acwm.co.la.ca.us/pdf/WhiteGardenSnaileng_pdf.
- Gentry, J. W. 1965. Crop insects of Northeast Africa-Southwest Asia (273). Agricultural Research Service; United States Department of Agriculture, Washington, D.C. 214 pp.
- Grund, R. 2002. South Australian Butterflies Datasheet: *Lampides boeticus* (Linnaeus) (Long-tailed Pea-blue). Chariot Pty Ltd. 4 pp.
- Hächler, M. 1986. Notes on three pests of subtropical ornamental plants captured in western Switzerland: *Trichoplusia orichalcea* Fab., *Syngrapha circumflexa* L. and *Spodoptera littoralis* Bois. (Lep., Noctuidae). / Notizen über drei in der Westschweiz gefangene subtropische Zierpflanzenschädlinge: *Trichoplusia orichalcea* Fab., *Syngrapha circumflexa* L. und *Spodoptera littoralis* Bois. (Lep., Noctuidae) [ABSTRACT]. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 59(3-4):263-266.
- Hammad, S. M. 1978. Pests of grain legumes and their control in Egypt. Pages 135-137 in S. R. Singh, H. F. Van Emden, and T. Ajibola Taylor, (eds.). *Pests of grain legumes: ecology and control*. Academic Press, London.
- Hanounik, S. B., and M. Bisri. 1991. Status of diseases of faba bean in the Mediterranean region and their control. *Options Méditerranéennes* 10:59-66.
- Harakly, F. A. 1975. Biological studies on the loopers *Autographa gamma* (L.) and *Cornutiplusia circumflexa* (L.) (Lep., Noctuidae) infesting truck crops in Egypt. *Zeitschrift für Angewandte Entomologie* 78(3):285-290.
- Harakly, F. A., and M. A. H. Assem. 1978. Ecological studies on the true pests of leguminous plants in Egypt. I. Biting and chewing pests. Pages 233-236 in *Proceedings of the Fourth Conference of Pest Control, September 30 - October 3, 1978. (Part I)*. Academy of Scientific Research and Technology and National Research Centre., Cairo; Egypt.
- Hashem, M. S., A. A. Abd-Elsamed, and A. A. A. Saleh. 2009. Monitoring and seasonal abundances of the leafhoppers; *Empoasca decipiens* (Paoli), *Empoasca decedens* (Paoli) and their associated predators on some leguminous vegetable crops in Egypt. *Egyptian Journal of Biological Pest Control* 19(2):105-114.

- Heu, R. 2007. Distribution and host records of agricultural pests and other organisms in Hawaii. State of Hawaii Department of Agriculture, Plant Industry Division, Plant Pest Control Branch, Survey Program, Honolulu, Hawaii. 71 pp.
- Hill, D. S. 1983. Agricultural insect pests of the tropics and their control. Cambridge University Press, Cambridge, U.S.A.
- Horton, D. 2008. Pink hibiscus mealybug found in north Georgia. The University of Georgia, College of Agricultural and Environmental Sciences.
- Ibrahim, I. K. A., Z. A. Handoo, and A. A. El-Sherbiny. 2000. A Survey of Phytoparasitic Nematodes on Cultivated and Non-Cultivated Plants in Northwestern Egypt. *Journal of Nematology* (Supplement) 32(4S):478-485.
- Iida, H., T. Kitamura, and K. Honda. 2009. Comparison of egg-hatching rate, survival rate and development time of the immature stage between B- and Q-biotypes of *Bemisia tabaci* (Gennadius) (Homoptera: Aleyrodidae) on various agricultural crops. *Applied Entomology and Zoology* 44(2):267-273.
- International Institute of Entomology. 1993. Distribution Maps of Pests: *Helicoverpa armigera* (Hübner) (Map No. 15 (2nd revision)). CAB International, London. 6 pp.
- IPPC. 2011. International Standards For Phytosanitary Measures, Publication No. 11: Pest Risk Analysis for Quarantine Pests Including Analysis of Environmental Risks and Living Modified Organisms. Food and Agriculture Organization of the United Nations, Secretariat of the International Plant Protection Convention (IPPC), Rome, Italy. 30 pp.
- IPPC. 2012. International Standards For Phytosanitary Measures, Publication No. 5: Glossary of Phytosanitary Terms. Food and Agriculture Organization of the United Nations, Secretariat of the International Plant Protection Convention (IPPC), Rome, Italy. 38 pp.
- Jacas, J. A., A. Hermosa de Mendoza, M. Cambra, and R. Balduque. 1997. *Asymmetrasca decedens* a new pest of almond in Spain (EPPO Bulletin 27). 523-524 pp.
- Janse, J. D., H. E. v. d. Beld, J. Elphinstone, S. Simpkins, N. N. A. Tjou-Tam-Sin, and J. Vaerenbergh, Van., 1994. Introduction to Europe of *Ralstonia solanacearum* biovar 2, race 3 in *Pelargonium zonale* cuttings. *Journal of Plant Pathol.* 86:147-155.
- Kaniuczak, Z. 2006. Occurrence and harmfulness of bean weevils (*Bruchus rufimanus* Boh.) in field bean. / Występowanie oraz szkodliwość strakowca bobowego (*Bruchus rufimanus* Boh.) na bobiku. *Progress in Plant Protection* 46(2):473-475.
- Karel, A. K., and A. Autrique. 1989. Insects and other pests in Africa. Pages 455-504 in H. F. Schwartz and M. A. Pastor-Corrales, (eds.). *Bean production problems in the tropics* (second edition). Centro Internacional de Agricultura Tropical (CIAT), Cali, Colombia.
- Kergoat, G. J., A. Delobel, and J.-F. Silvain. 2004. Phylogeny and host-specificity of European seed beetles (Coleoptera, Bruchidae), new insights from molecular and ecological data. *Molecular Phylogenetics and Evolution* 32(3):855-865.
- Khalafallah, E. M. E., I. A. Khodeir, and E. A. El-Srand. 2006. Survey of leafhopper species and their time of appearance in cotton fields at Kafr El-Shiekh [Abstract]. *Egypt. J. Agric. Res.* 84(4).
- Khalafallah, E. M. E., Z. Shenishen, I. S. El-Hawary, and M. A. Khattab. 2005. The host range and overwintering sites of *Nezara viridula* L. under Kafr El-Sheikh conditions. *Egyptian Journal of Agricultural Research* 83(1):79-85.
- Khodeir, I. A. 2006. Ecological notes on the leafhopper *Orosius albicinctus* distant (Homoptera: Cicadellidae) at Kafr El-Sheikh, Egypt [ABSTRACT]. *Bulletin of Faculty of Agriculture, Cairo University* 57(1):169-184.
- Koenning, S. R., Overstreet, C., , Noling, J.W., , P. A. Donald, J. O. Becker, and F. B.A. 1999. Survey of Crop Losses in Response to Phytoparasitic Nematodes in the United States for 1994. *Journal of Nematology* 31(587-618).
- Koike, S. T. 1998. Severe outbreak of chocolate spot of Fava bean, caused by *Botrytis fabae*, in California. *Disease Note. Plant Disease* 82(7):831.

- Lerner, R. 2001. Growing beans in the home vegetable garden. Purdue University Cooperative Extension Service, West Lafayette, IN. 3 pp.
- Liquido, N. J., R. T. Cunningham, and S. Nakagawa. 1990. Host Plants of Mediterranean Fruit Fly (Diptera: Tephritidae) on the island of Hawaii (1949-1985 Survey). *Journal of Economic Entomology* 83(5):1863-1878.
- Liquido, N. J., L. A. Shinoda, and R. T. Cunningham. 1991. Host plants of the Mediterranean fruit fly (Diptera: Tephritidae): An annotated world review. *Miscellaneous Publications of the Entomological Society of America* 77:1-52.
- Magarey, R., D. Borchert, and W. Schlegel. 2008. Global plant hardiness zones for phytosanitary risk analysis. *Scientia Agricola* 65(Special):54-59.
- Magsig-Castillo, J., J. G. Morses, G. P. Walker, J. L. Bi, P. F. Rugman-Jones, and R. Stouthamer. 2010. Phoretic Dispersal of Armored Scale Crawlers (Hemiptera: Diaspididae). *J. Econ. Entomol.* 103(4):1172-1179.
- Makkouk, K., L. Rizkallah, M. Madkour, M. El-Sherbeeney, S. Kumari, A. Amriti, and M. Solh. 1994. Survey of faba bean (*Vicia faba* L.) for viruses in Egypt *Phytopathol. Mediterr.* 33(3):20--211.
- Makkouk, K. M., S. G. Kumari, and L. Bos. 1990. Broad bean wilt virus: host range, purification, serology, transmission characteristics, and occurrence in faba bean in West Asia and North Africa. *Netherlands journal of plant pathology* 96(5):291-300.
- Mansour, M. H., A. E. Aboul-Nasr, and N. Y. Salem. 1981. The influence of different host plants on some biological aspects of *Heliothis armigera* (Hub.). *Experientia* 37(5):484-485.
- Mau, R. F. L., and J. L. M. Kessing. 1992. *Cyrtoblabe gnidiella* (Milliere). University of Hawaii, College of Tropical Agriculture and Human Resources, Hawaii Department of Agriculture. 2 pp.
- McGregor, B. M. 1987. Tropical Products Transport Handbook. USDA.
- Mckenzie, C. 2011. Q Biotype Whitefly Finds in North America. United States Department of Agriculture, Agricultural Research Service. 2 pp.
- Medjdoub-Bensaad, F., M. A. Khelil, and J. Huignard. 2007. Bioecology of broad bean bruchid *Bruchus rufimanus* Boh. (Coleoptera: Bruchidae) in a region of Kabylia in Algeria. *African Journal of Agricultural Research* 2(9):412-417.
- Membracoidea of the World Database. 2010. *Neolimnus egyptiacus* Matsumura 1908 (accepted name). Species2000 Catalogue of Life: Annual Checklist. <http://www.catalogueoflife.org/annual-checklist>. (Archived at PERAL).
- Migeon, A., and F. Dorkeld. 2006. Spider Mites Web: *Eutetranychus orientalis*. <http://www1.montpellier.inra.fr/CBGP/spmweb/notespecies.php?id=321>. (Archived at PERAL).
- Ministry of Agriculture and Land Reclamation. 2006. Green Beans. Ministry of Agriculture and Land Reclamation, Central Administration of Plant Quarantine (CAPQ) Cairo, Egypt.
- Ministry of Agriculture and Land Reclamation. 2011. Egypt green bean prerequisite request form (7CFR319.5). Ministry of Agriculture and Land Reclamation, Central Administration of Plant Quarantine (CAPQ) Cairo, Egypt. 3 pp.
- Muturi, J. J., J. P. Mbugi, J. M. Mueke, J. Lagerlof, J. K. Mungatu, G. Nyamasyo, and M. Gikungu. 2009. Collembola richness and diversity along a land-use intensity gradient in Taita, Kenya. *Tropical and Subtropical Agroecosystems* 11:415-422.
- Najar, A., S. Kumari, N. Attar, and S. Lababidi. 2011. First Report of Chickpea chlorotic stunt virus Infecting Legume Crops in Tunisia. *Plant Disease* 95(10):1321-1321.
- Nakhla, J. M., A. W. Tadros, and A. G. Hashem. 1997. Effect of fertilizers on the reduction of land snail population in citrus orchards. *Annals of Agricultural Science (Cairo)* 42(1):353-360.
- NASS. 2009. 2007 Census of Agriculture. United States Department of Agriculture, National Agricultural and Statistics Service (NASS). http://www.agcensus.usda.gov/Publications/2007/Online_Highlights/Desktop_Application/index.asp. (Archived at PERAL).
- National Invasive Species Council. No Date. Invasive Species of the Month: Pink Hibiscus Mealybug, *Maconellicoccus hirsutus*. National Invasive Species Council.

- Newberry, D. M., and M. G. Hill. 1985. Changes in the distribution of the coccid *Icerya seychellarum* Westw. on Aldabra Atoll in relation to vegetation density. Atoll Research Bulletin 291.
- Parihar, S.B.S., and O.P. Singh. 1988. Screening of certain potato varieties against lepidopterous pests. Indian Journal of Plant Protection, 16(1):83-85.
- PERAL. 2008. Importation of Fine Bean, *Phaseolus vulgaris* L., from Zambia into the Continental United States (Rev. 004). United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine, Center for Plant Health Science and Technology, Plant Epidemiology and Risk Analysis Laboratory, Raleigh, NC. 33 pp.
- PestID. 2012. Pest Identification Database (PestID). United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine. <https://mokes14.aphis.usda.gov/aqas/login.jsp>. (Archived at PERAL).
- PPQ. 2002. Electronic files for arthropods from pests not known to occur in the United States or of limited distribution and insects not known to occur in the United States. United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine (PPQ), Policy and Program Development, Risk Analysis Systems. 941 pp.
- PPQ. 2012. Guidelines for Plant Pest Risk Assessment of Imported Fruit and Vegetable Commodities (First Edition). United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine (PPQ), Raleigh, NC.
- Raupach, K., C. Borgemeister, M. Hommes, H.-M. Poehling, and M. Setamou. 2002. Effect of temperature and host plants on the bionomics of *Empoasca decipiens* (Homoptera: Cicadellidae). Crop Protection 21:113-119.
- Reed, W., and C.S. Pawar, 1982. *Heliothis*: a global problem. In: Reed W, Kumble V, ed. Proceedings of the International Workshop on Heliothis Management. ICRISAT Center, Patancheru, India, 15-20 November 1981 International Crops Research Institute for the Semi-Arid Tropics Patancheru, Andhra Pradesh India, 9-14.
- Robinson, G. S., P. R. Ackery, I. J. Kitching, G. W. Beccaloni, and L. M. Hernández. 2007a. HOSTS - a Database of the World's Lepidopteran Hostplants: *Spodoptera exempta*. London: Natural History Museum. (Archived at PERAL).
- Robinson, G. S., P. R. Ackery, I. J. Kitching, G. W. Beccaloni, and L. M. Hernández. 2007b. HOSTS - a Database of the World's Lepidopteran Hostplants: *Spodoptera littoralis*. London: Natural History Museum. (Archived at PERAL).
- Robinson, G. S., P. R. Ackery, I. J. Kitching, G. W. Beccaloni, and L. M. Hernández. 2011. HOSTS - a Database of the World's Lepidopteran Hostplants: *Chrysodeixis chalcites*. London: Natural History Museum. (Archived at PERAL).
- Rodríguez, I., H. Morales, J. M. Bueno, and C. Cardona M. No Date. The B biotype of *Bemisia tabaci* (Homoptera: Aleyrodidae) becomes more important in the Cauca Valley [Abstract]. Revista Colombiana de Entomología.
- Rungs, C. E. E. 1962. La faune nuisible aux tabacs [ABSTRACT]. Awamia 5:153-157.
- Schaefer, C. W., and A. R. Panizzi. 2000. Heteroptera of Economic Importance. CRC Press, Boca Raton. 828 pp.
- Seif, A., A. M. Varela, S. Michalik, and B. Lohr. 2001. A Guide to IPM in French Beans Production with Emphasis on Kenya. International Centre of Insect Physiology and Ecology (ICIPE) Science Press, Nairobi, Kenya.
- Shebl, M. A., S. M. Kamel, T. A. Abu Hashesh, and M. A. Osman. 2009. The most common insect species in Alfalfa fields in Egypt. Phegea 37(3):97-102.
- Shiba, T., K. Sugawara, and A. Arakawa. 2011. Evaluating the fungal endophyte *Neotyphodium occultans* for resistance to the rice leaf bug, *Trigonotylus caelestialium*, in Italian ryegrass, *Lolium multiflorum*. Entomologia Experimentalis et Applicata 141(1):45-51.
- Shishehbor, P., and P. A. Brennan. 1995. Parasitism of *Trialeurodes ricini* by *Encarsia formosa*: Level of parasitism, development time and mortality on different host plants. BioControl 40(3-4):299-305.

- Soliman, A. 2007. Egyptian crop technical files. Personal communication to Linda Logan USDA-APHIS Attache Cairo Egypt on September 5, 2007, from Dr. Ali Soliman, Head of Central Administration for Plant Quarantine.
- Spencer, K. A. 1965. A clarification of the status of *Liriomyza trifolii* (Burgess) and some related species (Diptera: Agromyzidae). *Proceedings of the Entomological Society of Washington* 67(1):32-40.
- Stang, D. 2009. Zipcode Zoo: *Maconellicoccus hirsutus*.
http://zipcodezoo.com/Animals/M/Maconellicoccus_hirsutus/ (Archived at PERAL).
- Sullivan, M., and E. Jones. 2010. Soybean Commodity-based Survey Guideline. United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine, Center for Plant Health Science and Technology, Cooperative Agriculture Pest Survey. 72 pp.
- Thindwa, H., and P. Khonje. 2005. Chapter 2.5: Malawi. Pages 150-156 in P. K. Anderson and F. J. Morales, (eds.). *Whitefly and Whitefly-Borne Viruses in the Tropics: Building a Knowledge Base for Global Action*. Centro Internacional de Agricultura Tropical, Cali, Colombia.
- Thomas, D. B., J. E. Eger, W. Jones, and G. Ortega-Leon. 2003. The African Cluster Bug, *Agonoscelis puberula* (Heteroptera: Pentatomidae), Established in the New World. *Florida Entomologist* 86(2):151-153.
- Thomas, M. C., J. B. Heppner, R. E. Woodruff, H. V. Weems, G. J. Steck, and T. R. Fasulo. 2005. Mediterranean Fruit Fly, *Ceratitis capitata* (Wiedemann) (Insecta: Diptera: Tephritidae) (EENY214). University of Florida, Institute of Food and Agricultural Sciences. 15 pp.
- Triplehorn, C. A., N. F. Johnson, and D. J. Borror. 2005. Borror and DeLong's introduction to the study of insects. Thompson Brooks/Cole.
- United States Fish and Wildlife Service. 2012. Endangered Species Database. United States Department of the Interior, United States Fish and Wildlife Service. <http://www.fws.gov/endangered/>. (Archived at PERAL).
- Unsinger, R. L. 1951. Heteroptera of the Marshall Islands. *Proceedings, Hawaiian Entomological Society* XIV(No. 2):315-321.
- van Huis, A., G. Woldewahid, K. Toleubayev, and W. van der Werf. 2008. Relationships between food quality and fitness in the desert locust, *Schistocerca gregaria*, and its distribution over habitats on the Red Sea coastal plain of Sudan. *Entomologia Experimentalis et Applicata* 127:144-156.
- Williams, D. J., and G. W. Watson. 1990. The Scale Insects of the Tropical South Pacific Region. Part 3. The Soft Scales (Coccidae) and Other Families. CAB International Institute of Entomology, London. 267 pp.
- Williams, M. C. 2008. Butterflies and Skippers of the Afrotropical Region (Papilionoidea and Hesperioidea) an Encyclopaedia File M - Lycaneinae, Polyommatainae (Lycaenidae) (Seventh edition), Rietondale, Pretoria. 155 pp.
- Zaher, M. A., K. K. Shehata, and H. El-Khatib. 1979. Population density effects on biology of *Tetranychus arabicus*, the common spider mite in Egypt [ABSTRACT]. Pages 507-509 in J. G. Rodriguez, ed. *Recent advances in acarology*. Volume I. Academic Press., New York; USA.
- Zhang, B. C. 1994. Index of Economically Important Lepidoptera. CAB International, Wallingford, UK. 599 pp.
- Zidan, Z. H., H.I. El-Deep, M. Wilson, and F.D.A. Asran. 1997. Molluscicidal activity of certain weed extracts and fertilizers against the brown garden snail, *Helix aspersa*. *Annals of Agricultural Science* 42(2):687-695.
- Zimmerman, E. C. 1958. Insects of Hawaii. A manual of the insects of the Hawaiian Islands, including an enumeration of the species and notes on their origin, distribution, hosts, parasites, etc. University of Hawaii Press, Honolulu. 542 pp.
- Zitter, T. A. 2005. Important New York Vegetable Diseases. Cornell University, Department of Plant Pathology. Last accessed June 11, 2012,
<http://vegetablemdonline.ppath.cornell.edu/NewsArticles/NYDiseases.htm>.

7. Appendix

Appendix A. Pests with non-actionable regulatory status

We found some evidence of the below listed organisms being associated with fresh green beans for consumption, in pods or shelled, and being present in the Egypt; however, because these organisms have non-actionable regulatory status for the continental United States, we did not list them in Table 2 of this risk assessment.

Below we list these organisms along with the references supporting their potential association with fresh green beans (*Phaseolus vulgaris* Linnaeus) for consumption, in pods or shelled, their potential presence in Egypt, their presence in the continental United States (if applicable), and their regulatory status for the continental United States. For organisms *not* present in the continental United States, we also provide justification for their non-actionable status.

Organism	Evidence and/or other notes
ARTHROPODS	
<i>Acanthoscelides obtectus</i> (Say)	PestID, 2012; CABI, 2012
<i>Acyrtosiphon pisum</i> (Harris) (= <i>Macrosiphum pisum</i> (Harris))	PestID, 2012; El Kifl et al., 1974; CABI, 2012
<i>Agrotis ipsilon</i> (Hufnagel)	El Kifl et al., 1974; PestID, 2012
<i>Aphis craccivora</i> Koch (= <i>A. laburni</i> Kaltenbach)	CABI, 2012; El Kifl et al., 1974; Favret, 2012; PestID, 2012
<i>Aphis fabae</i> Scopoli (= <i>A. compositae</i> Theobald)	PestID, 2012; CABI, 2012
<i>Aphis gossypii</i> Glover	CABI, 2012; PestID, 2012; CABI, 2012
<i>Aphis nerii</i> Boyer de Fonscolombe	CABI, 2012; Raupach et al., 2002; CABI, 2012
<i>Bemisia afer</i> (Priesner & Hosny)	Abd-Rabou and Ahmed, 2008; PestID, 2012; Thindwa and Khonje, 2005
<i>Bemisia tabaci</i> (Gennadius)	CABI, 2012; PestID, 2012; CABI, 2012
<i>Bemisia tabaci</i> (Gennadius) (B biotype) (= <i>B. argentifolii</i> Bellows, Perring, Gill and Hendrick) (CABI, 2012)	Abd-Rabou, 2006; CABI, 2012; PestID, 2012; Rodríguez et al., No Date; CABI, 2012
<i>Bemisia tabaci</i> (Gennadius) (Q biotype)	Brown, 2007; Iida et al., 2009; McKenzie, 2011; PestID, 2012
<i>Brevipalpus californicus</i> (Banks)	CABI, 2012; PestID, 2012; CABI, 2012
<i>Brevipalpus obovatus</i> Donnadieu	CABI, 2012; PestID, 2012; CABI, 2012
<i>Brevipalpus phoenicis</i> (Geijskes)	CABI, 2012; Childers et al., 2003; PestID, 2012; CABI, 2012
<i>Callosobruchus chinensis</i> (Linnaeus)	CABI, 2012; PestID, 2012; CABI, 2012
<i>Callosobruchus maculatus</i> (Fabricius)	Gentry, 1965; PestID, 2012; CABI, 2012
<i>Bruchus rufimanus</i> Boheman	Gentry, 1965; PestID, 2012; CABI, 2012
<i>Carpophilus hemipterus</i> (Linnaeus)	CABI, 2012; PestID, 2012; CABI, 2012
<i>Delia platura</i> (Meigen)	CABI, 2012
<i>Etiella zinckenella</i> (Treitschke)	PestID, 2012; CABI, 2012

Organism	Evidence and/or other notes
<i>Frankliniella schulzei</i> (Trybom)	CABI, 2012; PestID, 2012; Seif et al., 2001; CABI, 2012
<i>Gryllotalpa gryllotalpa</i> (Linnaeus)	CABI, 2012; El Kifl et al., 1974; PestID, 2012; CABI, 2012
<i>Hypera postica</i> (Gyllenhal)	PestID, 2012; CABI, 2012
<i>Lampides boeticus</i> Linnaeus	PestID, 2012; CABI, 2012
<i>Liriomyza trifolii</i> Burgess	PestID, 2012; CABI, 2012
<i>Macrosiphum euphorbiae</i> Thomas	PestID, 2012; CABI, 2012
<i>Myzus persicae</i> Sulzer	El-Lakwa et al., 1999; PestID, 2012
<i>Nezara viridula</i> (Linnaeus)	Khalafallah et al., 2005; PestID, 2012
<i>Ostrinia nubilalis</i> (Hübner)	CABI, 2012; PestID, 2012; CABI, 2012
<i>Pieris rapae</i> Linnaeus	El Kifl et al., 1974; PestID, 2012
<i>Pinnaspis buxi</i> (Bouché)	Ben-Dov et al., 2012; PestID, 2012
<i>Pinnaspis strachani</i> (Cooley)	Ben-Dov et al., 2012; PestID, 2012
<i>Planococcus citri</i> (Risso)	Ben-Dov et al., 2012; PestID, 2012
<i>Pseudaulacaspis pentagona</i> (Targoni-tozzetti)	Ben-Dov et al., 2012; PestID, 2012
<i>Rhopalosiphum rufiabdominale</i> (Sasaki)	PestID, 2012; CABI, 2012
<i>Sitona lineatus</i> Linnaeus	PestID, 2012; CABI, 2012
<i>Sitophilus zeamais</i> Motschulsky	PestID, 2012; CABI, 2012
<i>Smynthuroides betae</i> Westwood	Gentry, 1965; PestID, 2012
<i>Spodoptera exigua</i> (Hübner)	PestID, 2012; CABI, 2012
<i>Tetranychus cinnabarinus</i> (Boisduval) (= <i>T. cucurbitacearum</i> (Sayed), <i>T. telarius</i>)	PestID, 2012; CABI, 2012
<i>Tetranychus urticae</i> Koch (= <i>T. arabicus</i> Attiah)	PestID, 2012; Zaher et al., 1979; CABI, 2012
<i>Tetranychus atlanticus</i> McGregor	Canerday and Arant, 1964; Gentry, 1965; PestID, 2012
<i>Thrips tabaci</i> Lindeman	CABI, 2012; PestID, 2012; CABI, 2012
<i>Tribolium castaneum</i> Herbst	PestID, 2012; CABI, 2012
<i>Trichoplusia ni</i> (Hübner) (= <i>Plusia ni</i> (Hübner))	PestID, 2012; El Kifl et al., 1974; CABI, 2012
<i>Vanessa cardui</i> (Linnaeus)	Gentry, 1965; PestID, 2012
PATHOGENS	
Bacteria	
<i>Burkholderia cepacia</i> (Burkholder) Palleroni & Homes; (= <i>Pseudomonas cepacia</i> ; <i>Pseudomonas cepacia</i>)	EG, US: CABI, 2012; Bradbury, 1986
<i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i> (Jones) Hauben et al., comb. nov.; (= <i>Erwinia carotovora</i> subsp. <i>atroseptica</i> (Jones) Bergey et al.)	EG: CABI, 2012; US: Bradbury, 1986
<i>Pseudomonas marginalis</i> pv. <i>marginalis</i> (Brown) Stevens	EG: CABI, 2012; US: Bradbury, 1986,

Organism	Evidence and/or other notes
<i>Pseudomonas savastanoi</i> pv. <i>phaseolicola</i> (Burkholder) Gardan et al. (= <i>P. syringae</i> pv. <i>phaseolicola</i> (Burkholder) Young et al.)	EG: CABI, 2012; US: Bradbury, 1986
<i>Pseudomonas syringae</i> pv. <i>pisii</i> (Sackett) Young et al.	EG: US: CABI, 2012; US: Bradbury, 1986
<i>Pseudomonas syringae</i> pv. <i>syringae</i> van Hall	EG: CABI, 2012; US: Bradbury, 1986
<i>Pseudomonas viridiflava</i> (Burkholder) Dowson	EG: CABI, 2012; US: Bradbury, 1986
<i>Rhizobium radiobacter</i> (Beijerinck and van Delden) Young et al.	EG: CABI, 2012; US: Bradbury, 1986
<i>Xanthomonas axonopodis</i> pv. <i>alfalfae</i> (Riker et al.) Vauterin et al.	EG, US: CABI, 2012
<i>Xanthomonas axonopodis</i> pv. <i>phaseoli</i> (Smith) Vauterin et al. (= <i>X. campestris</i> pv. <i>phaseoli</i> (Smith) Dye)	EG: CABI, 2012; US: Bradbury, 1986
Fungi	
<i>Alternaria alternata</i> (Fr.) Keissl. (= <i>A. tenuis</i> Nees)	EG: Farr et al., 2007; Ministry of Agriculture and Land Reclamation, 2006
<i>Alternaria brassicicola</i> (Schwein.) Wiltshire	EG, US: CABI, 2012; Farr et al., 2007
<i>Aspergillus flavus</i> Link	EG, US: CABI, 2012; Farr et al., 2007
<i>Aspergillus niger</i> Tiegh.	EG: CABI, 2012; US: Farr et al., 2007
<i>Athelia rolfsii</i> (Curzi) Tu and Kimbr. (= <i>Corticium rolfsii</i> Curzi)	EG, US: CABI, 2012; Farr et al., 2007
<i>Botryotinia fuckeliana</i> (de Bary) Whetzel (= <i>Botrytis cinerea</i> Pers)	EG, US: Farr et al., 2007; CABI, 2012
<i>Botryosphaeria rhodina</i> (Berk. & M.A. Curtis) Arx (= <i>Lasiodiplodia theobromae</i> (Pat.) Griffon & Maubl.)	EG: Ministry of Agriculture and Land Reclamation, 2006; US: Farr et al., 2007
<i>Botrytis fabae</i> Sardiña	EG: Ministry of Agriculture and Land Reclamation, 2006; US: Farr et al., 2012; Koike, 1998; Zitter, 2005
<i>Choanephora cucurbitarum</i> (Berk. & Ravenel) Thaxt.	EG, US: CABI, 2012
<i>Cochliobolus lunatus</i> R.R. Nelson & Haasis	EG, US: CABI, 2012
<i>Cochliobolus sativus</i> (S. Ito & Kurib.) Drechsler ex Dastur (= <i>Helminthosporium sativum</i> Pammel, C.M. King and Bakke)	EG: CABI, 2012; US: Farr et al., 2007
<i>Colletotrichum truncatum</i> (Schwein.) Andrus & W.D. Moore	EG, US: CABI, 2012; US: Farr et al., 2007
<i>Corticium rolfsii</i> Curzi (= <i>Sclerotium rolfsii</i>)	EG, US: CABI, 2012
<i>Diaporthe phaseolorum</i> (Lehman) Wehm.	EG: CABI, 2012; US: Farr et al., 2007
<i>Erysiphe betae</i> (Vaňha) Weltzien (= <i>E. polygoni</i> DC.)	EG, US: CABI, 2012

Organism	Evidence and/or other notes
<i>Fusarium phaseoli</i> (Burkh.) T. Aoki & O'Donnell	EG, US: CABI, 2012
<i>Fusarium oxysporum</i> Schltdl.	EG: CABI, 2012; Ministry of Agriculture and Land Reclamation, 2006; US: Farr et al., 2007
<i>Glomerella cingulate</i> (Stoneman) Spauld. & H. Schrenk (= <i>Colletotrichum gloesporioides</i> (Penz.) Penz. and Sacc.; <i>Gibberella avenacea</i> R.J. Cook; <i>Fusarium avenaceum</i> (Fr.: Fr.) Sacc.)	EG, US: Farr et al., 2007
<i>Gibberella zeae</i> (Schwein.) Petch	EG: CABI, 2012; US: Farr et al., 2007
<i>Leveillula taurica</i> (Lév.) G. Arnaud	EG: CABI, 2012; US: Farr et al., 2007
<i>Mycosphaerella cruenta</i> Latham	EG: CABI, 2012; US: Farr et al., 2007
<i>Penicillium italicum</i> Wehmer	EG, US: CABI, 2012
<i>Phoma pinodella</i> (Jones) Morgan-Jones & Burch (= <i>Sphaerotheca fuliginea</i> (Schltdl.) Pollacci)	EG: CABI, 2012; US: Farr et al., 2007
<i>Phytophthora cryptogea</i> Pethybr. & Laff.	EG, US: CABI, 2012
<i>Podosphaera xanthii</i> (Castagne) U. Braun & Shishkoff (= <i>Sphaerotheca fuliginea</i> (schltdl.) Pollacci)	EG: CABI, 2012; US: Farr et al., 2007
<i>Pythium aphanidermatum</i> (Edson) Fitzp.	EG: Ministry of Agriculture and Land Reclamation, 2006; US: Farr et al., 2007
<i>Pythium debaryanum</i> Hesse	EG, US: Farr et al., 2007
<i>Pythium irregulare</i> Buisman	EG: EMOALR, 2006; US: Farr et al., 2007
<i>Pythium myriotylum</i> Drechsler	EG: CABI, 2012; US: Farr et al., 2007
<i>Pythium ultimum</i> Trow	EG, US: Farr et al., 2007
<i>Rosellinia necatrix</i> Berl. Ex	EG, US: Farr et al., 2007
<i>Sclerotinia sclerotiorum</i> (Lib.) de Bary	EG: Ministry of Agriculture and Land Reclamation, 2006; US: Farr et al., 2007
<i>Thanatephorus cucumeris</i> (A.B. Frank) Donk (= <i>Rhizoctonia solani</i> J.G. Kühn)	EG: Ministry of Agriculture and Land Reclamation, 2006; US: Farr et al., 2007; CABI, 2012
<i>Thielaviopsis basicola</i> (Berk. & Broome) Ferraris (= <i>Chalara elegans</i> Nag Raj & W.B. Kendr.)	EG: CABI, 2012; US: Farr et al., 2007
<i>Uromyces appendiculatus</i> F. Strauss (= <i>U. phaseoli</i> G. Winter)	EG: Ministry of Agriculture and Land Reclamation, 2006; US: Farr et al., 2007
<i>Uromyces viciae-fabae</i> (Pers.) J. Schröt.	EG: Ministry of Agriculture and Land Reclamation, 2006; US: Farr et al., 2007
<i>Verticillium dahliae</i> Kleb.	EG: CABI, 2012; US: Farr et al., 2007
Nematodes	
<i>Ditylenchus dipsaci</i> Kühn	EG: CABI, 2012, Hanounik and Bisri, 1991; US: US: Ferris, 2012
<i>Helicotylenchus dihystrera</i> (Cobb) Sher	EG, US: CABI, 2012
<i>Helicotylenchus multicinctus</i> (Cobb) Golden	EG, US: CABI, 2012;

Organism	Evidence and/or other notes
<i>Helicotylenchus pseudorobustus</i> (Steiner) Golden	EG, US: CABI, 2012
<i>Heterodera glycines</i> Ichinohe	EG, US: CABI, 2012
<i>Heterodera schachtii</i> A. Schdmit	EG, US: CABI, 2012; Ferris, 2012
<i>Meloidogyne arenaria</i> Neal Chitwood	EG: Abdel-Dali, 2007; Ibrahim et al., 2000; US: CABI, 2012
<i>Meloidogyne hapla</i> Chitwood	EG, US: CABI, 2012
<i>Meloidogyne incognita</i> (Kofoed and White) Chitwood	EG, US: CABI, 2012
<i>Pratylenchus brachyurus</i> (Godfrey) Filipjev and Schuurmans Stekhoven	EG, US: CABI, 2012
<i>Pratylenchus penetrans</i> (Cobb) Chitwood and Oteifa	EG, US: CABI, 2012
<i>Pratylenchus thornei</i> Sher and Allen	EG, US: CABI, 2012
<i>Pratylenchus vulnus</i> Allen& Jensen	EG: CABI, 2012; US: Koenning et al., 1999
Viruses	
<i>Alfalfa mosaic alfamovirus</i>	EG: CABI, 2012; Makkouk et al., 1994; US: CABI, 2012; , Brunt et al., 1996
<i>Bean common mosaic necrosis potyvirus</i> (BCMNV) (= <i>Bean common mosaic potyvirus</i> (BCMV))	EG: Makkouk et al., 1994; Makkouk et al., 1990; US: Brunt et al., 1996
<i>Bean leafroll luteovirus</i>	EG: Makkouk et al., 1994; US: CABI, 2012
<i>Bean yellow mosaic potyvirus</i> (BYMV)	EG: Makkouk et al., 1994; US: CABI, 2012
<i>Beet curly top curtovirus</i> (BCTV) (Geminiviridae)	EG: CABI, 2012; US: Creamer et al., 1996
<i>Broad bean wilt comovirus</i> (BBWV) (Fabaviridae)	EG, US: Brunt et al., 1996
<i>Chickpea chlorotic stunt virus</i>	EG: Najar et al., 2011 (listed in Tunisia but compared with EG isolates); US: Brunt et al., 1996
<i>Cucumber mosaic cucumovirus</i> (CMV)	EG: Makkouk et al., 1994; US: Brunt et al., 1996
<i>Peanut mottle potyvirus</i> (PeMoV)	EG: Brunt et al., 1996; US: Brunt et al., 1996
<i>Soybean dwarf luteovirus</i> (SbDV)	EG: Makkouk et al., 1994 ; US: (CA) Brunt et al., 1996
<i>Squash leaf curl virus begomovirus</i> (SCLCV) (= <i>Bean Calico Mosaic Virus</i>)	EG: CABI, 2012; US: Brunt et al., 1996
<i>Tobacco mosaic Virus tobamovirus</i> (TMV)	EG: Makkouk et al., 1994; US: Brunt et al., 1996
<i>Tobacco rattle tobnavirus</i> (TRV)	EG: CABI, 2012; US: Brunt et al., 1996
<i>Tobacco streak ilarvirus</i>	EG CABI, 2012; US: Brunt et al., 1996
<i>Tomato ringspot nepovirus</i>	EG: CABI, 2012; US: Brunt et al., 1996,
<i>Tomato spotted wilt tospovirus</i> (TSWV)	EG: Brunt et al., 1996; US: Brunt et al., 1996
<i>Tomato yellow leaf curl begomovirus</i> (TYLCV)	EG: Duffy and Holmes, 2007; EPPO, 2005; EPPO, 2012b
<i>Watermelon mosaic potyvirus</i> (WMV)	EG: CABI, 2012; US: Brunt et al., 1996